The nested $SU(N)$ off-shell Bethe Ansatz and exact form factors

H Babujian[1], A Foerster[2] and M Karowski[3]

[1] Yerevan Physics Institute, Alikhanian Brothers 2, Yerevan, 0036 Armenia
[2] Instituto de Física da UFRGS, Porto Alegre, RS - Brazil
[3] Institut für Theoretische Physik, Freie Universität Berlin, Arnimallee 14, 14195 Berlin, Germany

E-mail: babujian@physik.fu-berlin.de, angela@if.ufrgs.br, karowski@physik.fu-berlin.de

Abstract. The form factor equations are solved for an $SU(N)$ invariant S-matrix under the assumption that the anti-particle is identified with the bound state of $N-1$ particles. The solution is obtained explicitly in terms of the nested off-shell Bethe ansatz where the contribution from each level is written in terms of multiple contour integrals. The general solution is illustrated for some operators, such as the energy momentum tensor, the fields and the current.

PACS numbers: 11.10.-z; 11.10.Kk; 11.55.Ds

1. Introduction

Thirty-five years after the discovery of asymptotic freedom [1, 2], Quantum Chromodynamics (QCD), the theory of the strong interactions between quarks and gluons, remains a challenge. One of the most important trends in theoretical physics in the last several decades is the development of exact methods which are completely different from the perturbation theory since the perturbation theory is not applicable in the strong coupling regime. Resolution of the strong coupling problem would give us a full understanding of the structure of the interactions in non-Abelian gauge theory and in particular would shed light on the mysterious confinement phenomena in QCD. One promising possibility to overcome the limitations of the perturbation theory is the application of exact integrability. This is one of the reasons why investigation of two dimensional exactly solvable lattice statistical systems, 1-dimensional spin chains and quantum field theory models in $1+1$ dimensions in the framework of the Bethe ansatz method is still quite actual [3]. Some two dimensional exactly integrable field theories, such as nonlinear σ-model or Gross-Neveu model exhibit asymptotic freedom, a phenomenon initially observed in non-Abelian gauge theories. From this point of view the two-dimensional integrable quantum field theories are in a sense a laboratory for the investigations of those properties of quantum field theories,which can not be described via perturbation theory. One may hope that some of these properties, at least qualitatively, could be extended to the higher dimensions too.

The *Bethe ansatz method* [4], was first introduced by Bethe to solve the isotropic Heisenberg model. Yang [5] and Sutherland [6] generalized the technique of the Bethe ansatz for those cases where the underlying symmetry group is larger than $SU(2)$.

This method is now called the *nested Bethe ansatz*. A great impulse in the theory of integrable models was given by Faddeev and collaborators [7] with the development of the algebraic Bethe ansatz method. Another modification of the Bethe ansatz is the *off-shell Bethe ansatz*, which was originally formulated in [8] to calculate correlation function in WZNW models (see also [9, 10]). This version of the Bethe ansatz paves the way to an analysis of off-shell quantities and opens up the intriguing possibility to merge the Bethe ansatz and the form factor approach.

The exact determination of form factors, which are matrix elements of local operators, was formulated in the seventies [11]. This approach was developed further and studied in the context of several explicit models by Smirnov [12] . Subsequently [13] the techniques of the off-shell Bethe ansatz were used to determine the form factors for the sine-Gordon model. There, however, the underlying group structure is simple and there was no need to use a nested version of the off-shell Bethe ansatz. Recently the form factor program has received renewed interest in different areas, such as condensed matter physics [14, 15, 16] and atomic physics [17]. In particular, applications to Mott insulators and carbon nanotubes [18, 14] doped two-leg ladders [19] and in the field of Bose-Einstein condensates of ultracold atomic and molecular gases [20, 17] have been discussed. In addition, form factors have been also employed to compute corrections to the entanglement, an essential property in the field of Quantum Computation [21].

In the present article we will focus on the determination of the form factors for an $SU(N)$ invariant S-matrix. We have to apply the nested off-shell Bethe ansatz to get the vectorial part of the form factors. This solves the missing link of Smirnov's [12] formula for the $SU(N)$ form factors, where the vectors were given by an "indirect definition" characterized by necessary properties but not provided explicitly. The procedure is similar as for the scaling $Z(N)$ Ising and affine $A(N-1)$ Toda models [22, 23] because the bound state structures of these models are similar. However, the algebraic structure of the form factors for the $SU(N)$ model is more complicated, because the S-matrix possesses backward scattering. Therefore we have to develop a nontrivial algebraic *nested off-shell Bethe ansatz*. We note that $SU(N)$ form factors were also calculated in [12, 24, 25] using other techniques, see also the related paper [26, 27]. We believe that our integral representation may shed some light for a better understanding of the correlation functions of $SU(N)$ invariant models, which are also interesting for applications [28].

The paper is organized as follows: In Section 2 we recall some known formulae and derive some simple results, which will be used in the following. In Section 3 we construct the general form factor formula. In Section 4 we discuss form factors of some operators such as the energy momentum tensor, the Dirac field and the $SU(N)$ current. A summary of our results can be found in 5. In Appendix A we present some basic properties of the $SU(N)$ bound state S-matrix. In Appendix B we provide the proofs of the main theorem and in Appendix C we prove the lemma dedicated to the higher level Bethe ansatz.

2. General setting

2.1. The $SU(N)$ S-matrix

The general solutions of the Yang-Baxter equations, unitarity and crossing relations for a $U(N)$ invariant S-matrix have been obtained in [29]. The S-matrix for the scattering of two particles of rank 1 (belonging to the vector representation of $SU(N)$) can be written as

$$S_{\alpha\beta}^{\delta\gamma}(\theta) = \delta_{\alpha\gamma}\delta_{\beta\delta}\, b(\theta) + \delta_{\alpha\delta}\delta_{\beta\gamma}\, c(\theta)\,. \tag{1}$$

The amplitudes which we use in the following are given by

$$a(\theta) = b(\theta) + c(\theta) = -\frac{\Gamma\left(1-\frac{\theta}{2\pi i}\right)\Gamma\left(1-\frac{1}{N}+\frac{\theta}{2\pi i}\right)}{\Gamma\left(1+\frac{\theta}{2\pi i}\right)\Gamma\left(1-\frac{1}{N}-\frac{\theta}{2\pi i}\right)} \tag{2}$$

and

$$c(\theta) = -\frac{i\eta}{\theta} b(\theta)\,, \quad \eta = 2\pi/N.$$

The amplitude $a(\theta)$ is the highest weight S-matrix eigenvalue for the two particle scattering. It will be essential for the Bethe ansatz below.

The S-matrix eigenvalue $b(\theta) - c(\theta)$ has a pole at $\theta = i\eta$ which means that there exist bound states of r fundamental particles $\alpha_1 + \ldots + \alpha_r \to (\rho_1 \ldots \rho_r)$ (with $\rho_1 < \ldots < \rho_r$) which transform as the anti-symmetric $SU(N)$ tensor representation of rank r, $(0 < r < N)$. The masses of the bound states satisfy $m_r = m_{N-r}$ which suggests Swieca's [30, 31] picture that the antiparticle of a particle of rank r is to be identified with the particle of rank $N - r$ (see also [22, 23]).

The general bound state S-matrix formula [32, 33] for the scattering of a bound state with another particle can be used to calculate iteratively the scattering of a particle of rank $N-1$ with a particle of rank 1 (for details see Appendix A and [34]). The result is

$$S_{(\rho)\alpha}^{\gamma(\sigma)}(\theta) = (-1)^{N-1}\left(\delta_{(\rho)}^{(\sigma)}\delta_\alpha^\gamma\, b(\pi i-\theta) + \mathbf{C}^{\gamma(\sigma)}\mathbf{C}_{(\rho)\alpha}\, c(\pi i-\theta)\right) \tag{3}$$

where $(\sigma) = (\sigma_1 \ldots \sigma_{N-1})$ and $(\rho) = (\rho_1 \ldots \rho_{N-1})$ denote the bound states. We have introduced the charge conjugation matrices

$$\begin{aligned} &\mathbf{C}_{(\alpha_1\ldots\alpha_{N-1})\alpha_N} = \mathbf{C}_{\alpha_1(\alpha_2\ldots\alpha_N)} = \epsilon_{\alpha_1\ldots\alpha_N} \\ &\mathbf{C}^{\alpha_1(\alpha_2\ldots\alpha_N)} = \mathbf{C}^{(\alpha_1\ldots\alpha_{N-1})\alpha_N} = (-1)^{N-1}\epsilon^{\alpha_1\alpha_2\ldots\alpha_N} \end{aligned} \tag{4}$$

with $\epsilon_{\alpha_1\ldots\alpha_N}$ and $\epsilon^{\alpha_1\alpha_2\ldots\alpha_N}$ total anti-symmetric and $\epsilon_{1\ldots N} = \epsilon^{1\ldots N} = 1$. The charge conjugation matrices satisfy

$$\mathbf{C}_{\alpha(\rho)}\mathbf{C}^{(\rho)\beta} = \delta_\alpha^\beta \quad , \qquad \mathbf{C}_{\alpha(\rho)}A_\beta^\alpha\mathbf{C}^{\beta(\rho)} = (-1)^{N-1}\,\mathrm{tr}\,A \tag{5}$$

Formula (3) may be interpreted as an unusual (because of the sign $(-1)^{N-1}$) crossing relation

$$S_{(\rho)\alpha}^{\gamma(\sigma)}(\theta) = (-1)^{N-1}\mathbf{C}_{(\rho)\delta}S_{\alpha\beta}^{\delta\gamma}(\pi i-\theta)\mathbf{C}^{\beta(\sigma)}$$

supporting Swieca's picture that the antiparticle of a particle of rank 1 is to be identified with the bound state of rank $N-1$.

In order to simplify the formulae we extract the factor $a(\theta)$ from the S-matrix (1)

$$S_{\alpha\beta}^{\delta\gamma}(\theta) = a(\theta)\tilde{S}_{\alpha\beta}^{\delta\gamma}(\theta)$$

such that

$$\begin{aligned} &\tilde{S}_{\alpha\beta}^{\delta\gamma}(\theta) = \delta_\alpha^\gamma\delta_\beta^\delta\,\tilde{b}(\theta) + \delta_\alpha^\delta\delta_\beta^\gamma\,\tilde{c}(\theta) \\ &\tilde{b}(\theta) = \frac{\theta}{\theta-i\eta}\,, \quad \tilde{c}(\theta) = \frac{-i\eta}{\theta-i\eta}\,. \end{aligned} \tag{6}$$

2.2. Generalized Form factors

For a state of n particles of kind α_i with rapidities θ_i and a local operator $\mathcal{O}(x)$ we define the form factor functions $F^{\mathcal{O}}_{\alpha_1\dots\alpha_n}(\theta_1,\dots,\theta_n)$, or using a short hand notation $F^{\mathcal{O}}_{\underline{\alpha}}(\underline{\theta})$, by

$$\langle\, 0\,|\,\mathcal{O}(x)\,|\,\theta_1,\dots,\theta_n\,\rangle^{in}_{\underline{\alpha}} = e^{-ix(p_1+\dots+p_n)}F^{\mathcal{O}}_{\underline{\alpha}}(\underline{\theta}),\ \text{for } \theta_1>\ldots>\theta_n \quad (7)$$

where $\underline{\alpha}=(\alpha_1,\dots,\alpha_n)$ and $\underline{\theta}=(\theta_1,\dots,\theta_n)$. For all other arrangements of the rapidities the functions $F^{\mathcal{O}}_{\underline{\alpha}}(\underline{\theta})$ are given by analytic continuation. Note that the physical value of the form factor, i.e. the left hand side of (7), is given for ordered rapidities as indicated above and the statistics of the particles. The $F^{\mathcal{O}}_{\underline{\alpha}}(\underline{\theta})$ are considered as the components of a co-vector valued function $F^{\mathcal{O}}_{1\dots n}(\underline{\theta})\in V_{1\dots n}=\left(V^{1\dots n}\right)^{\dagger}$. The general form factor can be written as [11]

$$F^{\mathcal{O}}_{1\dots n}(\underline{\theta}) = K^{\mathcal{O}}_{1\dots n}(\underline{\theta})\prod_{1\le i<j\le n}F(\theta_{ij}) \quad (8)$$

where $F(\theta)$ is the minimal form factor function (9) and the K-function $K^{\mathcal{O}}_{1\dots n}(\underline{\theta})$ is given below. It satisfies the form factor equations (i) – (v) (see below and [11, 12, 13, 33, 23]). For bosons and fermions they follow from general LSZ-assumptions and "maximal analyticity" [13]. For anyons there are additional statistics factors [35, 23, 36]. As we will discuss below in section 3 the statistics factors are modified due to the unusual crossing relation of the S-matrix (see [31]). We will also provide a constructive and systematic way of how to solve the form factor equations (i) – (v) for the co-vector valued function $F^{\mathcal{O}}_{1\dots n}$ for the $SU(N)$ S-matrix.

Minimal form factor: The solutions of Watson's and the crossing equations (i) and (ii) for two particles with no poles in the physical strip $0\le\operatorname{Im}\theta\le\pi$ and at most a simple zero at $\theta=0$ are the minimal form factors. In particular those for highest weight states are essential for the construction of the off-shell Bethe ansatz. One easily finds the minimal solution of

$$F(\theta)=a(\theta)\,F(-\theta)=F(2\pi i-\theta)$$

using (2) as

$$F(\theta)=c\exp\int_0^\infty\frac{dt}{t\sinh^2 t}e^{\frac{t}{N}}\sinh t\,(1-1/N)\,(1-\cosh t\,(1-\theta/(i\pi)))\,. \quad (9)$$

It belongs to the highest weight vector $w=(2,0,\dots,0)$ of the two particle state. We define the corresponding 'Jost-function' as for the $Z(N)$ models [22, 23] by the equation

$$\prod_{k=0}^{N-2}\phi(\theta+ki\eta)\prod_{k=0}^{N-1}F(\theta+ki\eta)=1\ ,\quad \eta=2\pi/N \quad (10)$$

which is typical for models where the bound state of $N-1$ particles is the anti-particle [23]. The solution is

$$\phi(\theta)=\Gamma(\theta/(2\pi i))\,\Gamma(1-1/N-\theta/(2\pi i)) \quad (11)$$

and satisfies the relations

$$\begin{aligned}\phi(\theta) &= \phi(-\theta)a\,(-\theta) = \phi((N-1)i\eta - \theta)\\ &= \frac{1}{-b(\theta)}\phi(2\pi i - \theta) = \frac{a(\theta - 2\pi i)}{-b(\theta)}\phi(\theta - 2\pi i)\,. \end{aligned} \tag{12}$$

The equations (11) and (10) also determine the normalization constant c in (9) (for details see Appendix A and [34]).

2.3. Nested "off-shell" Bethe ansatz for $SU(N)$

We consider a state with n particles and define as usual in the context of the algebraic Bethe ansatz [7, 37] the monodromy matrix

$$\tilde T_{1\dots n,0}(\underline{\theta},\theta_0) = \tilde S_{10}(\theta_{10})\cdots \tilde S_{n0}(\theta_{n0}). \tag{13}$$

It is a matrix acting in the tensor product of the "quantum space" $V^{1\dots n} = V^1 \otimes \cdots \otimes V^n$ and the "auxiliary space" V^0. All vector spaces V^i are isomorphic to a space V whose basis vectors label all kinds of particles. Here we consider $V \cong \mathbb{C}^N$ as the space of the vector representation of $SU(N)$. The Yang-Baxter algebra relation for the S-matrix yields

$$\tilde T_{1\dots n,a}(\underline{\theta},\theta_a)\,\tilde T_{1\dots n,b}(\underline{\theta},\theta_b)\,\tilde S_{ab}(\theta_a-\theta_b) = \tilde S_{ab}(\theta_a-\theta_b)\,\tilde T_{1\dots n,b}(\underline{\theta},\theta_b)\,\tilde T_{1\dots n,a}(\underline{\theta},\theta_a)$$

which implies the basic algebraic properties of the sub-matrices A, B, C, D with respect to the auxiliary space defined by

$$\tilde T_{1\dots n,0}(\underline{\theta},z) \equiv \begin{pmatrix} \tilde A_{1\dots n}(\underline{\theta},z) & \tilde B_{1\dots n,\beta}(\underline{\theta},z)\\ \tilde C^{\beta}_{1\dots n}(\underline{\theta},z) & \tilde D^{\beta'}_{1\dots n,\beta}(\underline{\theta},z)\end{pmatrix} \tag{14}$$

where $2 \le \beta,\beta' \le N$.

An 'off-shell' Bethe ansatz for a co-vector valued function $K_{\underline{\alpha}}(\underline{\theta})$ is given by the multiple contour integral

$$\boxed{K_{\underline{\alpha}}(\underline{\theta}) = \frac{1}{m!}\int_{\mathcal{C}_{\underline{\theta}}}\frac{dz_1}{R}\cdots\int_{\mathcal{C}_{\underline{\theta}}}\frac{dz_m}{R}\,\tilde h(\underline{\theta},\underline{z})\,p(\underline{\theta},\underline{z})\,\tilde\Psi_{\underline{\alpha}}(\underline{\theta},\underline{z})} \tag{15}$$

with the scalar functions

$$\tilde h(\underline{\theta},\underline{z}) = \prod_{i=1}^{n}\prod_{j=1}^{m}\tilde\phi(\theta_i - z_j)\prod_{1\le i<j\le m}\tau(z_i-z_j) \tag{16}$$

$$\tau(z) = \frac{1}{\phi(\theta)\phi(-\theta)}\,,\quad \tilde\phi(\theta) = \phi(\theta)a(\theta) = \phi(-\theta)\,. \tag{17}$$

where the function $\phi(\theta)$ is defined by the equation (10) with the solution (11). Below we will use the co-vector valued function $K_{\underline{\alpha}}(\underline{\theta})$ to construct form factors for operators $\mathcal{O}(x)$. The scalar p-function $p(\underline{\theta},\underline{z})$ which is in general a simple function of e^{θ_i} and e^{z_j} (see below) will depend on the specific operator $\mathcal{O}(x)$. It turns out that by means of the integral representation we transform the complicated form factor equations (22) – (26) (which are in general matrix equations) to simple equations for the p-function (see section 3). The integration contour $\mathcal{C}_{\underline{\theta}}$ (see figure 1) can be characterized as follows: there is a finite number of complex numbers $a_i(\underline{\theta})$, $b_j(\underline{\theta})$ such that the positions of all poles of the integrand in (15) are of the form

$$(1):\ a_i(\underline{\theta}) + 2\pi i k\,,\quad k\in\mathbb{N}$$
$$(2):\ b_j(\underline{\theta}) - 2\pi i l\,,\quad l\in\mathbb{N}$$

then $\mathcal{C}_{\underline{\theta}}$ runs from $-\infty$ to $+\infty$ such that all poles (1) are above and all poles (2) are below the contour (this is usually not a straight line). This contour is just the same as that used for the definition of Meijer's G-function. The constant R is defined by $R = \oint_\theta dz \tilde{\phi}(\theta - z)$ where the integration contour is a small circle around $z = \theta$ as part of $\mathcal{C}_{\underline{\theta}}$. The K-function is in general a linear combination of the *fundamental building*



Figure 1. The integration contour $\mathcal{C}_{\underline{\theta}}$. The bullets refer to poles of the integrand in (15).

blocks [38, 22, 23] given by (15) - (17). We consider here cases where the sum consists only of one term.

Nesting: If in (14) the range of β's is non-trivial, i.e. if $N > 2$ the Bethe ansatz co-vectors are of the form

$$\tilde{\Psi}_{\underline{\alpha}}(\underline{\theta}, \underline{z}) = L_{\underline{\beta}}(\underline{z})\, \tilde{\Phi}_{\underline{\alpha}}^{\underline{\beta}}(\underline{\theta}, \underline{z}) \tag{18}$$

where summation over all $\underline{\beta} = (\beta_1, \ldots, \beta_m)$ with $\beta_i > 1$ is assumed. The basic Bethe ansatz co-vectors $\tilde{\Phi}_{1\ldots n}^{\underline{\beta}} \in \left(V^{1\ldots n}\right)^\dagger$ are defined as

$$\tilde{\Phi}_{1\ldots n}^{\underline{\beta}}(\underline{\theta}, \underline{z}) = \Omega_{1\ldots n} \tilde{C}_{1\ldots n}^{\beta_m}(\underline{\theta}, z_m) \cdots \tilde{C}_{1\ldots n}^{\beta_1}(\underline{\theta}, z_1). \tag{19}$$

Here the "pseudo-vacuum" is the highest weight co-vector (with weight vector $w = (n, 0, \ldots, 0)$)

$$\Omega_{1\ldots n} = e(1) \otimes \cdots \otimes e(1)$$

where the unit vectors $e(\alpha)$ $(\alpha = 1, \ldots, N)$ correspond to the particle of type α which belongs to the vector representation of $SU(N)$. The pseudo-vacuum vector satisfies

$$\begin{array}{rcl} \Omega_{1\ldots n}\, \tilde{B}_{1\ldots n}^{\beta}(\underline{\theta}, z) &=& 0 \\ \Omega_{1\ldots n}\, \tilde{A}_{1\ldots n}(\underline{\theta}, z) &=& \Omega_{1\ldots n} \\ \Omega_{1\ldots n}\, \tilde{D}_{1\ldots n,\beta}^{\beta'}(\underline{\theta}, z) &=& \delta_\beta^{\beta'} \prod_{i=1}^n \tilde{b}(\theta_i - z)\Omega_{1\ldots n}\,. \end{array} \tag{20}$$

The amplitudes of the scattering matrices are given by eqs. (1) and (2). The technique of the **'nested Bethe ansatz'** means that one makes for the coefficients $L_{\underline{\beta}}(\underline{z})$ in (18) the analogous construction (15) - (17) as for $K_{\underline{\alpha}}(\underline{\theta})$ where now the indices $\underline{\beta}$ take only the values $2 \le \beta_i \le N$. This nesting is repeated until the space of the coefficients becomes one dimensional. It is well known (see [39]) that the 'off-shell' Bethe ansatz states are highest weight states if they satisfy a matrix difference equation which is here the form factor equation (ii) (see below). The $SU(N)$ weights are

$$w = (n_0 - n_1, n_1 - n_2, \dots, n_{N-2} - n_{N-1}, n_{N-1}) \tag{21}$$

where $n_0 = n$, $n_1 = m, \dots$ are the numbers of variables in the various levels of the nesting.

3. $SU(N)$ form factors

In this section we perform the form factor program for the general $SU(N)$ S-matrix. For this purpose, we perform the nested Bethe ansatz method with $N-1$ levels combined with the off-shell Bethe ansatz. Applications of the results to the $SU(N)$ chiral Gross-Neveu model [40] will be investigated in a separate article [34]. We consider operators $\mathcal{O}(x)$ with charge $Q^{\mathcal{O}}$ where $Q^{\mathcal{O}}$ is the smallest number n of fundamental (rank 1) particles such that $\langle 0 \,|\, \mathcal{O}(x) \,|\, \theta_1, \dots, \theta_n \,\rangle^{in}$ is non-vanishing. For example for the energy momentum tensor and the $SU(N)$-current $Q^T = Q^J = 0$ and for the fundamental field $Q^\psi = 1$.

$SU(N)$ form factor equations: The co-vector valued auxiliary function $F^{\mathcal{O}}_{1\dots n}(\underline{\theta})$ is meromorphic in all variables $\theta_1, \dots, \theta_n$ and satisfies the following relations:

(i) The Watson's equations describe the symmetry property under the permutation of both, the variables θ_i, θ_j and the spaces $i, j = i+1$ at the same time

$$F^{\mathcal{O}}_{\dots ij \dots}(\dots, \theta_i, \theta_j, \dots) = F^{\mathcal{O}}_{\dots ji \dots}(\dots, \theta_j, \theta_i, \dots)\, S_{ij}(\theta_{ij}) \tag{22}$$

for all possible arrangements of the θ's.

(ii) The crossing relation implies a periodicity property under the cyclic permutation of the rapidity variables and spaces

$$\begin{aligned} {}^{\mathrm{out},\bar{1}}\langle\, \theta_1 \,|\, \mathcal{O}(0) \,|\, \theta_2, \dots, \theta_n \,\rangle^{\mathrm{in,conn.}}_{2\dots n} &= F^{\mathcal{O}}_{1\dots n}(\theta_1 + i\pi, \theta_2, \dots, \theta_n)\, \dot{\sigma}_1 \mathbf{C}^{\bar{1}1} \\ &= F^{\mathcal{O}}_{2\dots n1}(\theta_2, \dots, \theta_n, \theta_1 - i\pi) \mathbf{C}^{1\bar{1}} \end{aligned} \tag{23}$$

where $\dot{\sigma}^{\mathcal{O}}_\alpha$ takes into account the statistics of the particle α with respect to $\mathcal{O}$. The charge conjugation matrix $\mathbf{C}^{\bar{1}1}$ is defined by (4).

(iii) There are poles determined by one-particle states in each sub-channel given by a subset of particles of the state in (7). In particular the function $F^{\mathcal{O}}_{\underline{\alpha}}(\underline{\theta})$ has a pole at $\theta_{12} = i\pi$ such that

$$\operatorname*{Res}_{\theta_{12}=i\pi} F^{\mathcal{O}}_{1\dots n}(\theta_1, \dots, \theta_n) = 2i\, \mathbf{C}_{12}\, F^{\mathcal{O}}_{3\dots n}(\theta_3, \dots, \theta_n)\, (\mathbf{1} - \dot{\sigma}_2 S_{2n} \dots S_{23})\,. \tag{24}$$

(iv) If there are also bound states in the model the function $F^{\mathcal{O}}_{\underline{\alpha}}(\underline{\theta})$ has additional poles. If for instance the particles 1 and 2 form a bound state (12), there is a pole at $\theta_{12} = i\eta,\ (0 < \eta < \pi)$ such that

$$\operatorname*{Res}_{\theta_{12}=i\eta} F^{\mathcal{O}}_{12\dots n}(\theta_1, \theta_2, \dots, \theta_n) = F^{\mathcal{O}}_{(12)\dots n}(\theta_{(12)}, \dots, \theta_n)\, \sqrt{2}\Gamma^{(12)}_{12} \tag{25}$$

where the bound state intertwiner $\Gamma^{(12)}_{12}$ and the values of $\theta_1,\ \theta_2,\ \theta_{(12)}$ and η are given in [32, 33].

(v) Naturally, since we are dealing with relativistic quantum field theories we finally have

$$F^{\mathcal{O}}_{1\dots n}(\theta_1,\dots,\theta_n) = e^{s\mu}\,F^{\mathcal{O}}_{1\dots n}(\theta_1+\mu,\dots,\theta_n+\mu) \tag{26}$$

if the local operator transforms under Lorentz transformations as $\mathcal{O} \to e^{s\mu}\mathcal{O}$ where s is the "spin" of $\mathcal{O}$.

Here $\dot\sigma_1 = \rho\sigma^{\mathcal{O}}_1$ is a phase factor, where $\sigma^{\mathcal{O}}_1$ is the statistics factor of the operator $\mathcal{O}(x)$ with respect to the particle 1 and ρ is a sign factor due to the unusual crossing relation of the S-matrix which is determined as follows. The statistics factors in (ii) and (iii) are not arbitrary, consistency implies that both are the same and

$$\dot\sigma^{\mathcal{O}}_1\dot\sigma^{\mathcal{O}}_{\bar 1} = \left(\dot\sigma^{\mathcal{O}}_1\right)^N = (-1)^{(N-1)Q^{\mathcal{O}}}$$

with the solution

$$\sigma^{\mathcal{O}}_1 = e^{i\pi(1-1/N)Q^{\mathcal{O}}} \tag{27}$$

$$\rho = (-1)^{(N-1)+(1-1/N)\left(n-Q^{\mathcal{O}}\right)} = \pm 1 \tag{28}$$

where $Q^{\mathcal{O}} = n \,mod\, N$ is the charge of the operator $\mathcal{O}(x)$. This solution and the decomposition of $\dot\sigma^{\mathcal{O}}_1 = \rho\sigma^{\mathcal{O}}_1$ is of course not unique, the choice proposed is consistent with [31] and the usual relation of spin and statistics (for all examples discussed below).

3.1. The form factor formula and $SU(N)$ weights

In order to obtain a recursion relation where only form factors for the fundamental particles of type α (which transform as the $SU(N)$ vector representation) are involved, we have to apply iteratively the bound state relation (iv) to get the $(N-1)$-bound state which is to be identified with the anti-particle. Using then the annihilation residue equation (iii) we obtain with (5)

$$\operatorname*{Res}_{\theta_{N-1N}=i\eta}\dots\operatorname*{Res}_{\theta_{12}=i\eta} F^{\mathcal{O}}_{123\dots n}(\theta_1,\dots,\theta_n) \tag{29}$$
$$= 2i\sqrt{2}^{N-2}\Gamma\,\varepsilon_{1\dots N}F^{\mathcal{O}}_{N+1\dots n}(\theta_{N+1},\dots,\theta_n)\left(\mathbf{1}-\dot\sigma^{\mathcal{O}}_N S_{Nn}\dots S_{NN+1}\right)$$

where Γ is a constant (see (A.5) in Appendix A).

The properties of the K- function: We proposed above the n-particle form factors of an operator $\mathcal{O}(x)$ as given by the formula (8). In this expression the K-function $K^{\mathcal{O}}_{1\dots n}(\underline\theta)$ contains the entire pole structure and its properties are determined by the form factor equations (i) – (iii) which read in terms of the K-function as

$$K_{\dots ij\dots}(\dots,\theta_i,\theta_j,\dots) = K_{\dots ji\dots}(\dots,\theta_j,\theta_i,\dots)\,\tilde S_{ij}(\theta_{ij}) \tag{30}$$

$$K_{\alpha_1\alpha_2\dots\alpha_n}(\theta_1+i\pi,\theta_2,\dots,\theta_n)\,\dot\sigma_{\alpha_1}(-1)^{N-1} = K_{\alpha_2\dots\alpha_n\alpha_1}(\theta_2,\dots,\theta_n,\theta_1-i\pi) \tag{31}$$

$$\operatorname*{Res}_{\theta_{N-1N}=i\eta}\dots\operatorname*{Res}_{\theta_{12}=i\eta} K_{1\dots N}(\underline\theta) = c_0\prod_{i=N+1}^{n}\prod_{j=2}^{N}\tilde\phi(\theta_{ij})\,\varepsilon_{1\dots N}$$
$$\times K_{N+1\dots n}(\theta_{N+1},\dots,\theta_n)\left(\mathbf{1}-\dot\sigma_N S_{Nn}\dots S_{NN+1}\right) \tag{32}$$

with the constant

$$c_0 = 2i\sqrt{2}^{N-2}\Gamma\prod_{j=1}^{N-1}F^{-(N-j)}(ji\eta)\,. \tag{33}$$

For the second equation (31) the relation (5) for the charge conjugation matrices has been used and the third equation (32) follows from (10) and (29).

We propose the K-function $K_{\underline{\alpha}}(\underline{\theta})$ as the integral representation (15) with (16) and (17). The Bethe ansatz co-vector $\tilde{\Psi}_{\underline{\alpha}}(\underline{\theta},\underline{z})$ is of the form (18) - (19) and for the function $L_{\underline{\beta}}(\underline{z})$ one makes again an analogous ansatz as for the K-function (15) where the indices run over sets with one element less. For the $SU(N)$ case we have to iterate this $N-2$ times.

The higher level Bethe ansatz: We define for $l=1,\dots,N-2$, with the short notation $m=n_l$ and $k=n_{l+1}$

$$\begin{aligned} L^{(l)}_{\underline{\beta}}(\underline{z}) &= \frac{1}{k!}\int_{\mathcal{C}_{\underline{z}}}\frac{du_1}{R}\cdots\int_{\mathcal{C}_{\underline{z}}}\frac{du_k}{R}\,\tilde{h}(\underline{z},\underline{u})p^{(l)}(\underline{z},\underline{u})\,\tilde{\Psi}^{(l)}_{\underline{\beta}}(\underline{z},\underline{u}) \qquad (34)\\ \tilde{h}(\underline{z},\underline{u}) &= \prod_{i=1}^{m}\prod_{j=1}^{k}\tilde{\phi}(z_i-u_j)\prod_{1\le i<j\le k}\tau(u_i-u_j) \\ \tilde{\Psi}^{(l)}_{\underline{\beta}}(\underline{z},\underline{u}) &= L^{(l+1)}_{\underline{\gamma}}(\underline{u})\tilde{\Phi}^{(l-1)\underline{\gamma}}_{\underline{\beta}}(\underline{z},\underline{u}) \qquad (35)\end{aligned}$$

where $\underline{z}=z_1,\dots,z_m$, $\underline{u}=u_1,\dots,u_k$, $\underline{\beta}=\beta_1,\dots,\beta_m$, $\underline{\gamma}=\gamma_1,\dots,\gamma_k$. The indices assume the values $\beta_i=l+1,\dots,N$, $\gamma_i=l+2,\dots,N$. By this procedure one obtains the nested Bethe ansatz‡.

In order that the form factor $F^{\mathcal{O}}(\underline{\theta})$ given by (8), (15) - (19) and (34) satisfies the form factor equations (i) - (iii) the higher level L-functions $L^{(l)}_{\underline{\beta}}(\underline{z})$ have to satisfy:

(i)$^{(l)}$ Watson's equations

$$L^{(l)}_{\dots ij\dots}(\dots,z_i,z_j,\dots)=L^{(l)}_{\dots ji\dots}(\dots,z_j,z_i,\dots)\tilde{S}_{ij}(z_{ij})\,, \qquad (36)$$

(ii)$^{(l)}$ the crossing relation

$$L^{(l)}_{\beta_1,\beta_2,\dots,\beta_m}(z_1+2\pi i,z_2,\dots,z_m)=L^{(l)}_{\beta_2,\dots,\beta_m,\beta_1}(z_2,\dots,z_m,z_1) \qquad (37)$$

and

(iii)$^{(l)}$ the function $L^{(l)}_{\underline{\beta}}(\underline{z})$ has to possess simple poles at $z_{12},\dots,z_{N-l-1,N-l}=i\eta$ and it has to factorize in the neighborhood of these poles as

$$L^{(l)}_{\underline{\beta}}(\underline{z})\;\approx\;c_l\prod_{i=N-l+1}^{m}\prod_{j=2}^{N-l}\tilde{\phi}(z_{ij})\,\tilde{S}^{N\dots l+1}_{\underline{\hat{\beta}}}(\underline{\hat{z}})\,L^{(l)}_{\underline{\check{\beta}}}(\underline{\check{z}}) \qquad (38)$$

where we have used the short notations

$$\begin{aligned}\underline{\hat{\beta}}&=(\beta_1,\dots,\beta_{N-l})\,,\;\underline{\check{\beta}}=(\beta_{N-l+1},\dots,\beta_m)\\ \underline{\hat{z}}&=(z_1,\dots,z_{N-l}),\;\underline{\check{z}}=(z_{N-l+1},\dots,z_m).\end{aligned} \qquad (39)$$

The S-matrix $\tilde{S}^{N\dots l+1}_{\underline{\hat{\beta}}}(\underline{\hat{z}})$ describes the total scattering of the $N-l$ particles with rapidities $\underline{\hat{z}}$ for the initial and the final configuration of quantum numbers $\underline{\hat{\beta}}$ and $(N,\dots,l+1)$, respectively.

Theorem 1 *We make the following assumptions:*

‡ In [39] the nested off-shell Bethe ansatz was formulated in terms of "Jackson-type integrals" instead of contour integrals.

1. *the p-function $p(\underline{\theta},\underline{z})$ satisfies the equations*

$$\left.\begin{array}{ll}(\mathrm{i}') & p_{nm}(\underline{\theta},\underline{z}) \text{ is symmetric under } \theta_i \leftrightarrow \theta_j \text{ and } z_i \leftrightarrow z_j \\ (\mathrm{ii}'_1) & p_{nm}(\underline{\theta},\underline{z}) = \dot\sigma(-1)^{N-1+m} p_{nm}(\theta_1+2\pi i,\theta_2,\dots,\underline{z}) \\ (\mathrm{ii}'_2) & p_{nm}(\underline{\theta},\underline{z}) = (-1)^n p_{nm}(\underline{\theta},z_1+2\pi i,z_2,\dots) \\ (\mathrm{iii}'_1) & p_{nm}(\underline{\theta},\underline{\hat z}\underline{\check z}) = \nu\, p_{n-N\,m-N+1}(\underline{\check\theta},\underline{\check z}) \\ (\mathrm{iii}'_2) & p_{nm}(\underline{\theta},\underline{\hat z}\underline{\check z}) = (-1)^{m+N-1}\dot\sigma\,\nu\, p_{n-N\,m-N+1}(\underline{\check\theta},\underline{\check z})\end{array}\right\} \tag{40}$$

where in (iii'_1) : $\underline{\hat z} = (\theta_2,\dots,\theta_N)$ *and in* (iii'_2) : $\underline{\hat z} = (\theta_1,\dots,\theta_{N-1})$ *with* $\theta_{12} = \dots = \theta_{N-1N} = i\eta$. *In* (iii') *the short notations (39) and* $\underline{\check\theta} = (\theta_{N+1},\dots,\theta_n)$ *are used.*

2. *the higher level function $L^{(1)}_{\underline{\beta}}(\underline{z})$ satisfies* $(\mathrm{i})^{(1)}$ - $(\mathrm{iii})^{(1)}$ *of (36) – (38) and*
3. *the normalization constant ν satisfies*

$$\nu = \frac{2i\sqrt{2}^{N-2}\Gamma}{(N-1)!(i\eta)^{N-1}}\prod_{l=1}^{N-1}(-1)^{n_l+l}\left(\tilde\phi(li\eta)F(li\eta)\right)^{l-N} \tag{41}$$

then the co-vector valued function $F_{\underline{\alpha}}(\underline{\theta})$ given by the ansatz (8) and the integral representation (15) satisfies the form factor equations $(\mathrm{i}),(\mathrm{ii})$ *and* (iii) *in the form of (30) – (32).*

The proof of this theorem can be found in Appendix B.

Lemma 2 *The higher level function $L^{(l)}_{\underline{\beta}}(\underline{z})$ satisfies* $(\mathrm{i})^{(l)}$ - $(\mathrm{iii})^{(l)}$, *if the higher level p-function in the integral representation (34) satisfies*

$$\begin{array}{ll}(\mathrm{i}')^{(l)} & p^{(l)}(\underline{z},\underline{u}) \text{ is symmetric under } z_i \leftrightarrow z_j \\ (\mathrm{ii}')^{(l)} & \left\{\begin{array}{l} p^{(l)}_{mk}(\underline{z},\underline{u}) = (-1)^k p^{(l)}_{mk}(\underline{z}',\underline{u}) \\ \qquad\qquad\;\; = (-1)^m p^{(l)}_{mk}(\underline{z},\underline{u}')\end{array}\right. \\ (\mathrm{iii}')^{(l)} & p^{(l)}_{mk}(\underline{z},z_2,\dots z_{N-l},\underline{\check u})|_{z_{12}=\dots=z_{z_{N-l-1,N-l}}=i\eta} = p^{(l)}_{\check m\check k}(\underline{\check z},\underline{\check u})\end{array} \tag{42}$$

with $\check m = m-N+l$, $\check k = m-N+l-1$ *and* $m = n_l$, $k = n_{l+1}$. *We use the short notation* $\underline{z}' = z_1+2\pi i, z_2,\dots,z_m$ *and* $\underline{u}' = u_1+2\pi i,u_2,\dots,u_k$. *In* $(\mathrm{iii}')^{(l)}$ *the short notations (39) and* $\underline{\check u} = (u_{N-l},\dots,u_k)$ *are used.*

The constants satisfy the recursion relation

$$c_l = c_{l+1}(-1)^{n_{l+1}-N+l+1}\prod_{j=1}^{N-l-1}\tilde\phi(ji\eta) \tag{43}$$

with the solution

$$c_1 = \tilde\phi^{N-2}(i\eta)\tilde\phi^{N-3}(2i\eta)\cdots\tilde\phi((N-2)i\eta)\prod_{l=1}^{N-2}(-1)^{n_{l+1}-N+l+1} \tag{44}$$

if $c_{N-1} = 1$

This lemma is proved in Appendix C.

A compact form for full form factors: We may write the above results as

$$K_{\underline{\alpha}}(\underline{\theta}) = \frac{1}{\underline{n}!}\int \underline{\underline{dz}}\,\tilde{h}(\underline{\theta},\underline{\underline{z}})\,p(\underline{\theta},\underline{\underline{z}})\,\tilde{\Psi}_{\underline{\alpha}}(\underline{\theta},\underline{\underline{z}}) \tag{45}$$

$$\tilde{h}(\underline{\theta},\underline{\underline{z}}) = \tilde{h}(\underline{\theta},\underline{z}^{(1)})\cdots\tilde{h}(\underline{z}^{(N-2)},\underline{z}^{(N-1)}) \tag{46}$$

$$\tilde{\Psi}_{\underline{\alpha}}(\underline{\theta},\underline{\underline{z}}) = \tilde{\Phi}_{\underline{\beta}^{(N-1)}}(\underline{z}^{(N-2)},\underline{z}^{(N-1)})\cdots\tilde{\Phi}_{\underline{\alpha}}^{\underline{\beta}^{(1)}}(\underline{\theta},\underline{z}^{(1)}) \tag{47}$$

with the short notations

$$\int\underline{\underline{dz}} = \int\underline{dz}^{(1)}\cdots\int\underline{dz}^{(N-1)};\ \int\underline{dz}^{(l)} = \int_{\mathcal{C}_{\underline{z}^{(l-1)}}}\frac{dz_1^{(l)}}{R}\cdots\int_{\mathcal{C}_{\underline{z}^{(l-1)}}}\frac{dz_{n_l}^{(l)}}{R}$$

$$\underline{n}! = n_1!\cdots n_{N-1}!;\ \underline{\underline{z}} = \underline{z}^{(1)},\ldots,\underline{z}^{(N-1)};\ \underline{z}^{(l)} = z_1^{(l)},\ldots,z_{n_l}^{(l)}.$$

The function $\tilde{h}(\underline{\theta},\underline{\underline{z}})$ is given by (16) and (17).

The p-function and $SU(N)$ weights: In general the function $p(\underline{\theta},\underline{\underline{z}})$ depends on the rapidities $\underline{\theta}$ and all integration variables $\underline{z}^{(l)}$ $l=1,\ldots,N-1$. In order that the form factor $F^{\mathcal{O}}(\theta)$ given by (8) and (45) satisfies the form factor equations (i) - (iii) the p-function $p^{\mathcal{O}}(\underline{\theta},\underline{\underline{z}})$ in (15) which depends on the explicit nature of the local operator $\mathcal{O}(x)$ is assumed to satisfy

$$\left.\begin{array}{ll}
(\mathrm{i}') & p(\underline{\theta},\underline{\underline{z}}) \text{ is symmetric under } \theta_i\leftrightarrow\theta_j \text{ and } z_i^{(l)}\leftrightarrow z_j^{(l)}\\
(\mathrm{ii}_1') & p(\underline{\theta},\underline{\underline{z}}) = \dot{\sigma}(-1)^{n_1+N-1}p(\theta_1+2\pi i,\theta_2,\ldots,\underline{\underline{z}})\\
(\mathrm{ii}_2') & p(\underline{\theta},\underline{\underline{z}}) = (-1)^{w_l^{\mathcal{O}}+w_{l+1}^{\mathcal{O}}}p(\underline{\theta},\ldots,z_i^{(l)}+2\pi i,\ldots),\ l=1,\ldots,N-1\\
(\mathrm{iii}_1') & p(\underline{\theta},\underline{\hat{z}}^{(1)}\underline{\check{z}}^{(1)},\ldots) = \nu\,p(\underline{\check{\theta}},\underline{\check{z}}^{(1)},\ldots)\\
(\mathrm{iii}_2') & p(\underline{\theta},\underline{\hat{z}}^{(1)}\underline{\check{z}}^{(1)},\ldots) = (-1)^{n_1+N-1}\dot{\sigma}\,\nu\,p(\underline{\check{\theta}},\underline{\check{z}}^{(1)},\ldots)\\
(\mathrm{iii}_3') & p(\underline{\theta},\ldots,\underline{z}^{(l)},\underline{\hat{z}}^{(l+1)}\underline{\check{z}}^{(l+1)},\ldots) = p(\underline{\theta},\ldots,\underline{z}^{(l)},\underline{\check{z}}^{(l+1)},\ldots)\\
(\mathrm{v}') & p(\underline{\theta},\underline{\underline{z}}) = e^{s\mu}\,p(\underline{\theta}+\mu,\underline{\underline{z}}+\mu)
\end{array}\right\} \tag{48}$$

where in (iii_1') : $\underline{\hat{z}}^{(1)} = (\theta_2,\ldots,\theta_N)$ in (iii_2') : $\underline{\hat{z}}^{(1)} = (\theta_1,\ldots,\theta_{N-1})$ and in (iii_3') : $\underline{\hat{z}}^{(l+1)} = (z_2^{(l)},\ldots z_{N-l}^{(l)})$, furthermore in (iii_1') and (iii_2') : $\theta_{12}=\ldots=\theta_{N-1N}=i\eta$ and in (iii_3') : $z_{12}^{(l)}=\ldots=z_{z_{N-l-1,N-l}}^{(l)}=i\eta$. Here we have also included the equation (v′) which implies the behaviour of the form factor under Lorentz transformations (v) of (26). The sign factor (ii_2') follows from (40) and (42) as $(-1)^{n_{l-1}-n_{l+1}}$. This sign does not depend on the specific state of $F_{\underline{\alpha}}^{\mathcal{O}}$ but only on the weights of the operator $\mathcal{O}(x)$. This can be seen as follows: We assume that the operator $\mathcal{O}(x)$ transforms under $SU(N)$ as a highest weight representation with highest weight vector

$$w^{\mathcal{O}} = \left(w_1^{\mathcal{O}},\ldots,w_N^{\mathcal{O}}\right). \tag{49}$$

Then $SU(N)$ invariance implies that the weight vector of the form factor $F_{\underline{\alpha}}(\underline{\theta})$ as co-vector is of the form

$$w = w^{\mathcal{O}} + L(1,\ldots,1),\quad (L=0,1,\ldots) \tag{50}$$

because $w=(1,\ldots,1)$ correspond to the vacuum sector. Then because of (21)

$$(-1)^{n_{l-1}-n_{l+1}} = (-1)^{w_l+w_{l+1}} = (-1)^{w_l^{\mathcal{O}}+w_{l+1}^{\mathcal{O}}+2L} = (-1)^{w_l^{\mathcal{O}}+w_{l+1}^{\mathcal{O}}}.$$

Similarly, also the phase factor in (ii′) and (iii′) of (48) $\dot{\sigma}(-1)^{n_1+N-1}$ with $\dot{\sigma}=\rho\sigma$ and

$$\rho(-1)^{n_1+N-1} = (-1)^{(1-1/N)\left(n_0-Q^{\mathcal{O}}\right)+n_1} \tag{51}$$

does not depend on the specific state of $F_{\underline{\alpha}}^{\mathcal{O}}$ but only on the weights of the operator (see (28)).

4. Examples

To illustrate our general results we present some simple examples of solutions of the equations (48) for the p-functions. Details of the following calculations will be published elsewhere [34].

4.1. The energy momentum tensor

The local operator $\mathcal{O}(x) = T^{\rho\sigma}(x)$ (where $\rho, \sigma = \pm$ denote the light cone components) has charge $Q^T = 0$, is bosonic and transforms as a scalar under $SU(N)$ and has the weight vector of (49) $w^T = (0, \ldots, 0)$. The p-function (as for the sine-Gordon model in [33])

$$p^{T^{\rho\sigma}}(\underline{\theta}, \underline{z}) = N_n^{T^{\rho\sigma}} \sum_{i=1}^{n} e^{\rho\theta_i} \sum_{i=1}^{m} e^{\sigma z_i} .$$

solves (48) with

charge	$Q^T = 0$
weight vector	$w^T = (0, \ldots, 0)$
statistics factor	$\sigma^T = 1$
spin	$s^T = 2$

if $N_n^{T^{\rho\sigma}} / N_{n-N}^{T^{\rho\sigma}} = \nu$. The sign factor (51) is here $\rho(-1)^{n_1+N-1} = 1$.

For the $n = N$ particle form factor there are $n_l = N - l$ integrations in the l-th level of the off-shell Bethe ansatz because (see (21) and (50))

$$w = (n - n_1, n_1 - n_2, \ldots, n_{N-2} - n_{N-1}, n_{N-1}) = (1, 1, \ldots, 1, 1) .$$

We calculate the form factor of the particle α and the bound state $(\underline{\beta}) = (\beta_1, \ldots, \beta_{N-1})$ of $N-1$ particles. In each level all integrations up to one may be performed iteratively using the bound state relation (iv) (similar as in the proof of theorem 1). The result for the form factor of the particle α and the bound state $(\underline{\beta})$ writes as

$$\begin{aligned} F^{T^{\rho\sigma}}_{\alpha(\underline{\beta})}(\theta_1, \theta_2) &= K^{T^{\rho\sigma}}_{\alpha(\underline{\beta})}(\theta_1, \theta_2)\, G(\theta_{12}) \\ K^{T^{\rho\sigma}}_{\alpha(\underline{\beta})}(\theta_1, \theta_2) &= N_N^{T^{\rho\sigma}} \left(e^{\rho\theta_1} + e^{\rho\theta_2}\right) \int_{\mathcal{C}_{\underline{\theta}}} \frac{dz}{R} \tilde{\phi}(\theta_1 - z) e^{\sigma z} L(\theta_2 - z) \\ &\quad \times \epsilon_{\delta\underline{\gamma}} \tilde{S}^{\delta 1}_{\alpha\epsilon}(\theta_1 - z) \tilde{S}^{\epsilon(\underline{\gamma})}_{(\underline{\beta})1}(\theta_2 - z) \end{aligned} \tag{52}$$

where the summation is over $\underline{\gamma}$ and $\delta > 1$ and $G(\theta)$ is the minimal form factor function of two particles of rank $r = 1$ and $r = N - 1$. The functions $G(\theta)$ and $L(\theta)$ are given by

$$\begin{aligned} &G(i\pi - \theta) F(\theta) \phi(\theta) = 1 \\ &L(\theta) = \Gamma\left(\tfrac{1}{2} + \theta/(2\pi i)\right) \Gamma\left(-\tfrac{1}{2} + 1/N - \theta/(2\pi i)\right) . \end{aligned}$$

The remaining integral in (52) may be performed (similar as in [33]) with the result§

$$\langle\, 0 \,|\, T^{\rho\sigma}(0) \,|\, \theta_1, \theta_2 \,\rangle^{in}_{\alpha\underline{\beta})} = 4M^2 \epsilon_{\alpha\underline{\beta}} e^{\frac{1}{2}(\rho+\sigma)(\theta_1+\theta_2+i\pi)} \frac{\sinh\frac{1}{2}\left(\theta_{12} - i\pi\right)}{\theta_{12} - i\pi} G(\theta_{12}).$$

Similar as in [33] one can prove the eigenvalue equation

$$\left(\int dx T^{\pm 0}(x) - \sum_{i=1}^{n} p_i^{\pm}\right) |\, \theta_1, \ldots, \theta_n \rangle^{in}_{\underline{\alpha}} = 0$$

for arbitrary states.

§ In [41, 39] this result has been obtained using Jackson type integrals.

4.2. The fields $\psi_\alpha^{(\pm)}(x)$

Because the Bethe ansatz yields highest weight states we obtain the matrix elements of the spinor field $\psi(x) = \psi_1^{(+)}(x)$, where we consider here the + component of the spinor. The weight vector is $w^\psi = (1, 0, \ldots, 0)$ and the charge is $Q^\psi = 1$. The p-function for the local operator $\psi(x)$ is (see also [13])

$$p^\psi(\underline{\theta}, \underline{z}) = N_n^\psi \exp \frac{1}{2} \left((1 - 1/N) \sum_{i=1}^{n} \theta_i - \sum_{i=1}^{m} z_i \right).$$

It solves (48) with

charge	$Q^\psi = 1$
weight vector	$w^\psi = (1, \ldots, 0)$
statistics factor	$\sigma^\psi = e^{i\pi(1-1/N)}$
spin	$s^\psi = \frac{1}{2}(1 - 1/N)$

if $N_n^\psi / N_{n-N}^\psi = \nu$. The last two formulae are consistent with the proposal of Swieca et al. [30, 31] that the statistics of the fundamental particles in the chiral $SU(N)$ Gross-Neveu model should be $\sigma = \exp(2\pi i s)$, where $s = \frac{1}{2}\left(1 - \frac{1}{N}\right)$ is the spin (see also (27)). The sign factor (51) is here again $\rho(-1)^{n_1+N-1} = 1$.

The 1-particle form factor is for example

$$\langle 0 \,|\, \psi(0) \,|\, \theta \rangle_\alpha = \delta_{\alpha 1}\, e^{\frac{1}{2}(1-1/N)\theta}.$$

For the $n = N + 1$ particle form factor there are again $n_l = N - l$ integrations in the l-th level of the off-shell Bethe ansatz and the $SU(N)$ weights are $w = (2, 1, \ldots, 1, 1)$. Similar as above one obtains the 2-particle and 1-bound state form factor

$$F^\psi_{\alpha\beta(\underline{\gamma})}(\theta_1, \theta_2, \theta_3) = K^\psi_{\alpha\beta(\underline{\gamma})}(\theta_1, \theta_2, \theta_3) F(\theta_{12}) G(\theta_{13}) G(\theta_{23}),$$

$$\begin{aligned} K^\psi_{\alpha\beta(\underline{\gamma})} = N^\psi e^{\frac{1}{2}(1-1/N)\sum \theta_i} \int_{\mathcal{C}_{\underline{\theta}}} \frac{dz}{R} \tilde{\phi}(\theta_1 - z) \tilde{\phi}(\theta_2 - z) L(\theta_3 - z) e^{-\frac{1}{2}z} \\ \times \epsilon_{\delta\underline{\gamma}} \tilde{S}^{\delta 1}_{\alpha_1 \epsilon}(\theta_1 - z) \tilde{S}^{\epsilon 1}_{\alpha_2 \zeta}(\theta_2 - z) \tilde{S}^{\zeta(\underline{\gamma})}_{(\underline{\beta})1}(\theta_3 - z). \end{aligned}$$

We were not able to perform this integration. In [34] we will discuss the $1/N$ expansion of this exact form factor and compare the result with the $1/N$ expansion for the chiral Gross-Neveu model.

4.3. The current $J^\mu_{\alpha(\rho)}(x)$

The $SU(N)$ current transforms as the adjoint representation with highest weights $w^J = (2, 1, \ldots, 1, 0)$ and has charge $Q_J = 0$. Again, because the Bethe ansatz yields highest weight states we obtain the matrix elements of the highest weight component

$$\begin{aligned} J^\mu_{\alpha(\rho)} &= \delta_{\alpha 1} \epsilon_{(\rho)N} J^\mu, \quad (\rho) = (\rho_1, \ldots, \rho_{N-1}) \\ J^\mu &= \epsilon^{\mu\nu} \partial_\nu \varphi \end{aligned}$$

with the pseudo potential $\varphi(x)$. The p-function for the operator $\varphi(x)$

$$p^\varphi(\underline{\theta}, \underline{z}, \underline{u}) = N_n^\varphi \left(\sum_{i=1}^{n} \exp \theta_i \right)^{-1} \exp \frac{1}{2} \left(\sum_{i=1}^{n_0} \theta_i - \sum_{i=1}^{n_1} z_i^{(1)} - \sum_{i=1}^{n_{N-1}} z_i^{(N-1)} \right)$$

solves (48) with

charge	$Q^\varphi = 0$
weight vector	$w^\varphi = (2, 1, \ldots, 1, 0)$
statistics factor	$\sigma^\varphi = 1$
spin	$s^\varphi = 0,\ s^J = 1$

if $N_n^\varphi / N_{n-N}^\varphi = \nu$. The sign factor (51) is here $\rho(-1)^{n_1+N-1} = -1$.

We calculate the form factor of the particle α and the bound state $(\lambda) = (\lambda_1, \ldots, \lambda_{N-1})$ of $N-1$ particles with weight vector $w = (2, 1, \ldots, 1, 0)$. In each level the integrations may be performed iteratively using the bound state relation (iv), similar as above, however, here no integration remains. The result is

$$\begin{aligned} F^\varphi_{\alpha(\lambda)}(\theta,\omega) &= K^\varphi_{\alpha(\lambda)}(\theta,\omega)\, G(\theta-\omega) \\ K^\varphi_{\alpha(\lambda)}(\theta,\omega) &= N_2^\varphi \delta_{\alpha 1} \epsilon_{(\lambda)N} \frac{e^{\frac{1}{2}(\theta+\omega)}}{e^\theta + e^\omega}\,. \end{aligned}$$

The current form factor is therefore

$$\begin{aligned} F^{J^\pm_{\beta(\rho)}}_{\alpha(\lambda)}(\theta,\omega) &= \langle\, 0 \,|\, J^\pm_{\beta(\rho)}(0) \,|\, \theta,\omega \,\rangle^{in}_{\alpha(\lambda)} = \pm N_2 \delta^\beta_\alpha \delta^{(\rho)}_{(\lambda)} \left(e^{\pm\theta} + e^{\pm\omega}\right) \frac{e^{\frac{1}{2}(\theta+\omega)}}{e^\theta + e^\omega} G(\theta-\omega) \\ &= \delta^\beta_\alpha \delta^{(\rho)}_{(\lambda)}\, \bar v(\omega) \gamma^\pm u(\theta)\, G(\theta-\omega)/G(i\pi) \end{aligned}$$

as expected.

5. Summary

In this paper we have obtained the form factors for an $SU(N)$ invariant S-matrix. Since the S-matrix also describes backward scattering, the algebraic structure of the form factors was more intricate compared to the $Z(N)$ case [23]. We had to combine the nesting procedure with the techniques of the off-shell Bethe ansatz to capture the vectorial nature of the form factors. The solution was obtained explicitly in terms of multiple contour integrals. We exemplified our general solution for several operators, such as the energy momentum tensor, the fields and the current. We believe that our new integral representation may shed some light for a better comprehension of complicated objects, such as the correlation functions for the $SU(N)$ invariant Gross-Neveu model.

Acknowledgments

We thank R. Schrader, B. Schroer, and A. Zapletal for useful discussions. H.B. was supported partially by the Humboldt Foundation. A.F. acknowledges support from DAAD (Deutscher Akademischer Austausch Dienst), CAPES (Coordenação de Aperfeiçoamento Pessoal de Nivel Superior) and CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico). This work is also supported by the EU network EUCLID, 'Integrable models and applications: from strings to condensed matter', HPRN-CT-2002-00325.

Appendix A. The $SU(N)$ bound state S-matrix

The eigenvalue $S_-(\theta)$ has a pole at $\theta = i\eta = 2\pi i/N$ which means that there exist bound states of r fundamental particles $\alpha_1 + \ldots + \alpha_r \;\to\; (\rho_1 \ldots \rho_r)$ (with

$\rho_1 < \ldots < \rho_r$) which transform as the anti-symmetric $SU(N)$ tensor representation of rank r, $(0 < r < N)$. The masses of the bound states satisfy $m_r = m_{N-r}$ which suggests Swieca's [30, 42, 31] picture that the antiparticle of a particle of rank r is to be identified with the particle of rank $N-r$ (see also [22, 23]).

The general bound state S-matrix formula [32, 33] for the scattering of a bound state with another particle reads as

$$S_{(\rho\sigma)\gamma}^{\gamma'(\rho'\sigma')}(\theta_{(12)3})\Gamma_{\alpha\beta}^{(\rho\sigma)} = \Gamma_{\alpha'\beta'}^{(\rho'\sigma')} S_{\alpha\gamma''}^{\gamma'\alpha'}(\theta_{13}) S_{\beta\gamma}^{\gamma''\beta'}(\theta_{23}) \Big|_{\theta_{12}=i\eta} \tag{A.1}$$

where $\theta_{(12)}$ is the bound state rapidity and η the bound state fusion angle. The bound state fusion intertwiner $\Gamma_{\alpha\beta}^{(\rho\sigma)}$ is defined by

$$i \operatorname*{Res}_{\theta=i\eta} S_{\alpha\beta}^{\beta'\alpha'}(\theta) = \sum_{\rho<\sigma} \Gamma_{(\rho\sigma)}^{\beta'\alpha'} \Gamma_{\alpha\beta}^{(\rho\sigma)}. \tag{A.2}$$

With a convenient choice of an undetermined phase factor one obtains

$$\Gamma_{\alpha\beta}^{(\rho\sigma)} = \Gamma_{(\rho\sigma)}^{\beta\alpha} = i\sqrt{ia(i\eta)i\eta}\left(\delta_\alpha^\rho \delta_\beta^\sigma - \delta_\alpha^\sigma \delta_\beta^\rho\right) . \tag{A.3}$$

Iterating the general bound state formula $N-1$ times one obtains for the scattering of a bound state $(\rho) = (\rho_1\rho_2 \ldots \rho_{N-1})$ with another particle δ

$$S_{(\rho)\delta}^{\delta'(\sigma)}(\theta)\Gamma_{\underline{\alpha}}^{(\rho)} = \Gamma_{\underline{\alpha}'}^{(\sigma)} S_{\alpha_1\delta_1}^{\delta'\alpha_1'}(\theta + i\pi - i\eta) \ldots S_{\alpha_{N-1}\delta}^{\delta_{N-2}\alpha_{N-1}'}(\theta - i\pi + i\eta)$$

with the total bound state fusion intertwiner

$$\Gamma_{\underline{\alpha}}^{(\rho)} = \Gamma_{\alpha_1\alpha_2\ldots\alpha_{N-1}}^{(\rho_1\rho_2\ldots\rho_{N-1})} = \Gamma_{(\rho_1\rho_2\ldots\rho_{N-2})\alpha_{N-1}}^{(\rho_1\rho_2\ldots\rho_{N-1})} \cdots \Gamma_{(\rho_1\rho_2)\alpha_3}^{(\rho_1\rho_2\rho_3)} \Gamma_{\alpha_1\alpha_2}^{(\rho_1\rho_2)} .$$

Taking special cases for the external particles we obtain

$$S_{(1,2\ldots,N-1)N}^{N(1,2\ldots,N-1)}(\theta) = b(\theta + i\pi - i\eta) \ldots b(\theta - i\pi + i\eta) = (-1)^{N-1} a(\pi i - \theta)$$

$$S_{(1,2\ldots,N-1)N-1}^{N-1(1,2\ldots,N-1)}(\theta) = b(\theta + i\pi - i\eta) \ldots a(\theta - i\pi + i\eta) = (-1)^{N-1} b(\pi i - \theta)$$

$$S_{(2,3\ldots,N)1}^{N(1,2\ldots,N-1)}(\theta) = (-1)^N b(\theta + i\pi - i\eta) \ldots c(\theta - i\pi + i\eta) = c(\pi i - \theta) .$$

These results may be interpreted as an unusual crossing relation

$$\begin{aligned} \mathbf{C}_{(\beta)\beta}\Gamma_{\underline{\beta}}^{(\beta)} \left(S(\theta_1) \ldots S(\theta_{N-1})\right)_{\underline{\alpha}\delta}^{\delta'\underline{\beta}} &= \mathbf{C}_{(\beta)\beta} S_{(\alpha)\delta}^{\delta'(\beta)}(\theta)\Gamma_{\underline{\alpha}}^{(\alpha)} \\ &= (-1)^{N-1}\mathbf{C}_{(\alpha)\gamma}\Gamma_{\underline{\alpha}}^{(\alpha)}\, S_{\delta\beta}^{\gamma\delta'}(i\pi - \theta) \end{aligned} \tag{A.4}$$

with $\theta_j = \theta + i\pi - ji\eta$ and the charge conjugation matrix $\mathbf{C}_{(\beta)\beta}$ defined by (4). The total bound state fusion intertwiner satisfies

$$\mathbf{C}_{(\beta_1\ldots\beta_{N-1})\gamma}\Gamma_{\alpha_1\ldots\alpha_{N-1}}^{(\beta_1\ldots\beta_{N-1})} = \epsilon_{\alpha_1\ldots\alpha_{N-1}\gamma}\Gamma \;, \quad \Gamma = i\sqrt{\frac{1}{2\pi} N \Gamma^N \left(1 - 1/N\right)} . \tag{A.5}$$

We consider the total N-particle S-matrix (consisting of $N(N-1)/2$ factors) in terms of $\tilde{S}$

$$\tilde{S}_{12\ldots N} = \left(\tilde{S}_{12}\tilde{S}_{13}\ldots\tilde{S}_{1N}\right)\left(\tilde{S}_{23}\ldots\tilde{S}_{2N}\right)\ldots\tilde{S}_{N-1N} \tag{A.6}$$

in the limit $\theta_{jj+1} \to i\eta$ $(j = 1, \ldots N-1)$. It behaves as

$$\tilde{S}_{\alpha_1\alpha_2\ldots\alpha_N}^{\beta_N\ldots\beta_2\beta_1} \approx (N-1)! \frac{i\eta}{\theta_{12} - i\eta} \cdots \frac{i\eta}{\theta_{N-1N} - i\eta}\, \epsilon^{\beta_1\ldots\beta_N}\epsilon_{\alpha_1\ldots\alpha_N} . \tag{A.7}$$

The algebraic structure of this relation follows because one can use the Yang-Baxter equations to shift in (A.6) any factor $\tilde{S}_{ii+1} \sim (1-\mathbf{P})_{ii+1}$ to the right or the left. Therefore the expression is totally anti-symmetric with respect to the α_i and the β_i. The factor follows from (6).

The normalization constant c in (9) is calculated from (11) and (10) as

$$c = \Gamma^{-2(1-1/N)}\left(\frac{1}{2}-\frac{1}{2N}\right)\exp\left(-\int_0^\infty e^{\frac{1}{N}t}\left(\frac{\sinh\left(1-\frac{1}{N}\right)t}{t\sinh^2 t}-\frac{\left(1-\frac{1}{N}\right)}{t\sinh t}\right)dt\right) > 0\,.$$

Appendix B. Proof of Theorem 1

Proof. Property (i) in the form of (30) follows directly from (i$'$) of (40), the Yang-Baxter equations and the action of the S-matrix on the pseudo-ground state Ω

$$\begin{aligned}\Omega_{\dots ji\dots}\,\tilde{C}_{\dots ji\dots}(\dots\theta_j,\theta_i\dots)\,\tilde{S}_{ij}(\theta_{ij}) &= \Omega_{\dots ji\dots}\,\tilde{S}_{ij}(\theta_{ij})\,\tilde{C}_{\dots ij\dots}(\dots\theta_i,\theta_j\dots)\\ &= \Omega_{\dots ij\dots}\,\tilde{C}_{\dots ij\dots}(\dots\theta_i,\theta_j\dots)\,.\end{aligned}$$

because $\tilde{S}_{11}^{11}(\theta) = S_{11}^{11}(\theta)/a(\theta) = 1$ and $F(\theta) = F(-\theta)a(\theta)$.

Using (i) and (5) the property (ii) in the form of (31) may be rewritten as a matrix difference equation [43, 41, 39]

$$K_{1\dots n}(\underline{\theta})Q_{1\dots n}(\underline{\theta},i) = (-1)^{N-1}K_{1\dots n}(\underline{\theta}')\dot{\sigma}_1 \tag{B.1}$$

where $\underline{\theta}' = (\theta_1,\dots,\theta_i+2\pi i,\dots,\theta_n)$ and $\dot{\sigma}$ is a statistics factor. The matrix $Q(\underline{\theta},i)$ is the trace with respect to the auxiliary space

$$Q_{1\dots n}(\underline{\theta},i) = \mathrm{tr}_0\tilde{T}_{Q,1\dots n,0}(\underline{\theta},i) \tag{B.2}$$

of a modified monodromy matrix

$$\tilde{T}_{Q,1\dots n,0}(\underline{\theta},i) = \tilde{S}_{10}(\theta_1-\theta_i')\cdots\mathbf{P}_{i0}\cdots\tilde{S}_{n0}(\theta_n-\theta_i)$$

where $\mathbf{P} = \tilde{S}(0)$ is the permutation matrix. In (B.2) we use the rule that the rapidity of a line changes by $2\pi i$ if the line bends by 360^0 in the positive sense. Because of (i) it is sufficient to proof the matrix difference equation (B.1) for $i=1$, i.e. for $Q(\underline{\theta}) = Q(\underline{\theta},1)$.

In the following we will suppress the indices $1\dots n$. The Yang-Baxter relations (14) imply the typical commutation rules for the matrices $\tilde{A},\tilde{C},\tilde{D}$ defined in (14)

$$\tilde{C}^\beta(\underline{\theta},z)\tilde{A}(\underline{\theta},\theta) = \frac{1}{\tilde{b}(\theta-z)}\tilde{A}(\underline{\theta},\theta)\tilde{C}^\beta(\underline{\theta},z) - \frac{\tilde{c}(\theta-z)}{\tilde{b}(\theta-z)}\tilde{A}(\underline{\theta},z)\tilde{C}^\beta(\underline{\theta},\theta) \tag{B.3}$$

$$\begin{aligned}\tilde{C}^\beta(\underline{\theta},z)\tilde{D}_\gamma^{\gamma'}(\underline{\theta},\theta) &= \frac{1}{\tilde{b}(z-\theta)}\tilde{S}_{\beta'\gamma''}^{\gamma'\beta}(z-\theta)\tilde{D}_\gamma^{\gamma''}(\underline{\theta},\theta)\tilde{C}^{\beta'}(\underline{\theta},z)\\ &\quad -\frac{\tilde{c}(z-\theta)}{\tilde{b}(z-\theta)}\tilde{D}_\gamma^\beta(\underline{\theta},z)\tilde{C}^{\gamma'}(\underline{\theta},\theta)\end{aligned}$$

where $\beta,\beta',\gamma,\gamma',\gamma'' \in \{2,\dots,N\}$. In addition there are the Zapletal commutation rules [43, 41, 39] where also the matrices A_Q, C_Q, D_Q defined by

$$\tilde{T}_Q(\underline{\theta}) = \begin{pmatrix}\tilde{A}_Q(\underline{\theta}) & \tilde{B}_Q^\beta(\underline{\theta})\\ \tilde{C}_Q^\beta(\underline{\theta}) & \tilde{D}_{Q\,\beta}^{\;\beta'}(\underline{\theta})\end{pmatrix}$$

are involved [43]

$$\tilde{C}^{\beta}(\underline{\theta},z)\tilde{A}_Q(\underline{\theta}) = \frac{1}{\tilde{b}(\theta_1'-z)}\tilde{A}_Q(\underline{\theta})\tilde{C}^{\beta}(\underline{\theta}',z) - \frac{\tilde{c}(\theta_1'-z)}{\tilde{b}(\theta_1'-z)}\tilde{A}(\underline{\theta},z)\tilde{C}_Q^{\beta}(\underline{\theta}) \tag{B.4}$$

$$\begin{aligned}\tilde{C}^{\beta}(\underline{\theta},z)\tilde{D}_{Q\gamma}^{\gamma'}(\underline{\theta}) &= \frac{1}{\tilde{b}(z-\theta_1')}\tilde{S}_{\beta'\gamma''}^{\gamma'\beta}(z-\theta_1')\tilde{D}_{Q\gamma}^{\gamma''}(\underline{\theta})\tilde{C}^{\beta'}(\underline{\theta}',z)\\ &-\frac{\tilde{c}(z-\theta_1)}{\tilde{b}(z-\theta_1)}\tilde{D}_{\gamma}^{\beta}(\underline{\theta},z)\tilde{C}_Q^{\gamma'}(\underline{\theta})\,.\end{aligned} \tag{B.5}$$

Note that we assign to the auxiliary space of $\tilde{T}_Q(\underline{\theta})$ corresponding to the horizontal line the spectral parameter θ_1 on the right hand side and $\theta_1' = \theta_1 + 2\pi i$ on the left hand side.

We are now going to prove (B.1) in the form

$$K(\underline{\theta})\,Q(\underline{\theta}) = K(\underline{\theta})\left(\tilde{A}_Q(\underline{\theta}) + \sum_{\beta=2}^{N}\tilde{D}_{Q\beta}^{\beta}(\underline{\theta})\right) = \dot{\sigma}(-1)^{N-1}K(\underline{\theta}') \tag{B.6}$$

where $K(\underline{\theta})$ is a co-vector valued function as given by (15) and the Bethe ansatz state (18) and (19). Using a short notation we write

$$K(\underline{\theta}) = \int_{\mathcal{C}_{\underline{\theta}}} d\underline{z}\,\tilde{h}(\underline{\theta},\underline{z})p(\underline{\theta},\underline{z})\,\tilde{\Psi}(\underline{\theta},\underline{z})$$

(with $\int_{\mathcal{C}_{\underline{\theta}}} d\underline{z} = \frac{N_n}{m!}\int_{\mathcal{C}_{\underline{\theta}}}\frac{dz_1}{R}\cdots\int_{\mathcal{C}_{\underline{\theta}}}\frac{dz_m}{R}$). To analyze the left hand side of (B.6) we proceed as follows: We apply the trace of $\tilde{T}_Q$ to the co-vector $\tilde{\Psi}(\underline{\theta},\underline{z})$.

In the contribution from $\tilde{A}_Q(\underline{\theta})$ which means $\gamma' = \gamma = 1$ one may use Yang-Baxter relations to observe that only the amplitudes $\tilde{S}_{11}^{11}(\theta_1 - z_j) = 1$ appear in the S-matrices $\tilde{S}(\theta_1 - z_j)$ which are constituents of the C-operators. Therefore we may shift all z_j-integration contours $\mathcal{C}_{\underline{\theta}}$ to $\mathcal{C}_{\underline{\theta}'}$ without changing the values of the integrals, because there are no singularities inside $\mathcal{C}_{\underline{\theta}} \cup -\mathcal{C}_{\underline{\theta}'}$ (c.f. figure 1).

We now proceed as usual in the algebraic Bethe ansatz and push the $\tilde{A}_Q(\underline{\theta})$ and $\tilde{D}_Q(\underline{\theta})$ through all the $\tilde{C}$-operators using the commutation rules (B.4) and (B.5) and obtain

$$\begin{aligned}\tilde{C}^{\beta_m}(\underline{\theta},z_m)\cdots\tilde{C}^{\beta_1}(\underline{\theta},z_1)\,\tilde{A}_Q(\underline{\theta}) &= \prod_{j=1}^{m}\frac{1}{\tilde{b}(\theta_1'-z_j)}\,\tilde{A}_Q(\underline{\theta})\\ &\times\tilde{C}^{\beta_m}(\underline{\theta}',z_m)\cdots\tilde{C}^{\beta_1}(\underline{\theta}',z_1) + \sum uw_A,\end{aligned}$$

$$\begin{aligned}\tilde{C}^{\beta_m}(\underline{\theta},z_m)\cdots\tilde{C}^{\beta_1}(\underline{\theta},z_1)\,\tilde{D}_{Q\beta}^{\beta'}(\underline{\theta}) &= \prod_{j=1}^{m}\frac{1}{\tilde{b}(z_j-\theta_1')}\tilde{T}^{(1)}{}_{\underline{\beta}'\,\beta''}^{\beta'\underline{\beta}}(\underline{z},\theta_1')\tilde{D}_{Q\beta}^{\beta''}(\underline{\theta})\\ &\times\tilde{C}^{\beta_m'}(\underline{\theta},z_m)\cdots\tilde{C}^{\beta_1'}(\underline{\theta},z_1) + \sum uw_D\,.\end{aligned}$$

The "wanted terms" written out explicitly originate from the first term in the commutations rules (B.4); all other contributions yield the so-called "unwanted terms". The next level monodromy matrix is

$$\tilde{T}^{(1)}{}_{\underline{\beta}'\,\beta}^{\beta'\underline{\beta}}(\underline{z},\theta) = \left(\tilde{S}_{10}(z_1-\theta)\cdots\tilde{S}_{m0}(z_m-\theta)\right)_{\underline{\beta}'\,\beta}^{\beta'\,\underline{\beta}}$$

where the β's and also the internal summation indices take the values $2, \dots, N$. If we insert these equations into the representation (15) of $K(\underline{\theta})$ we first find that the wanted contribution from $\tilde{A}_Q$ already gives the result we are looking for. Secondly the wanted contribution from $\tilde{D}_Q$ applied to Ω gives zero. Thirdly the unwanted contributions from $\tilde{A}_Q$ and $\tilde{D}_Q$ cancel after integration over the z_j. These three facts follow from the 'shift relations' of the ϕ-function (12) and of the higher level L-function (ii)$^{(1)}$ of (37).

The proof of (iii) is similar to that for the $Z(N)$ model in [23]. We use the short hand notations

$$\begin{aligned}
&\underline{\theta} = (\theta_1, \dots, \theta_n),\ \hat{\underline{\theta}} = (\theta_1, \dots, \theta_N),\ \check{\underline{\theta}} = (\theta_{N+1}, \dots, \theta_n),\\
&\underline{\alpha} = (\alpha_1, \dots, \alpha_n),\ \hat{\underline{\alpha}} = (\alpha_1, \dots, \alpha_N),\ \check{\underline{\alpha}} = (\alpha_{N+1}, \dots, \alpha_n),\\
&\underline{z} = (z_1, \dots, z_m),\ \hat{\underline{z}} = (z_1, \dots, z_{N-1}),\ \check{\underline{z}} = (z_N, \dots, z_m),\\
&\underline{\beta} = (\beta_1, \dots, \beta_m),\ \hat{\underline{\beta}} = (\beta_1, \dots, \beta_{N-1}),\ \check{\underline{\beta}} = (\beta_N, \dots, \beta_m)\,.
\end{aligned}$$

We prove the form factor equation in the form of (32)

$$\begin{aligned}
\operatorname*{Res}_{\theta_{N-1N}=i\eta} \dots \operatorname*{Res}_{\theta_{12}=i\eta} K^{\mathcal{O}}_{1\dots N}(\underline{\theta}) &= c_0 \prod_{i=N+1}^{n} \prod_{j=2}^{N} \tilde{\phi}(\theta_{ij})\, \varepsilon_{1\dots N}\\
&\times K^{\mathcal{O}}_{N+1\dots n}(\theta_{N+1}, \dots, \theta_n) \left(\mathbf{1} - \dot{\sigma}^{\mathcal{O}}_N S_{Nn} \dots S_{NN+1}\right) \qquad \text{(B.7)}
\end{aligned}$$

for the K-function. The residues of $K_{1\dots n}(\underline{\theta})$ consists of three terms

$$Res_{\theta_{N-1N}=i\eta} \dots Res_{\theta_{12}=i\eta} K_{1\dots n}(\underline{\theta}) = R^{(1)}_{1\dots n} + R^{(2)}_{1\dots n} + R^{(3)}_{1\dots n}$$

because $N-1$ of the z integration contours will be "pinched" at three points. Again due to symmetry it is sufficient to determine the contribution from the $z_1, \dots, z_{N-1}$-integrations and multiply the result by $m \dots (m-N+2)$. The pinching points are

(1) $z_1 = \theta_2, \dots, z_{N-1} = \theta_N$,

(2) $z_1 = \theta_1, \dots, z_{N-1} = \theta_{N-1}$,

(3) $z_1 = \theta_2 - i\eta, \dots, z_{N-1} = \theta_N - i\eta$,

The contribution of (1) is given by $N-1$ integrations along small circles around $z_1 = \theta_2, z_2 = \theta_3, \dots, z_{N-1} = \theta_N$ (see figure 1). The S-matrices $\tilde{S}(\theta_2 - z_1), \dots, \tilde{S}(\theta_N - z_{N-1})$ yield the permutation operator $\tilde{S}(0) = \mathbf{P}$. Therefore for $\theta_{12}, \dots, \theta_{N-2N-1}, \theta_{N-1N} \to i\eta$ and $z_1 = \theta_2, z_2 = \theta_3, \dots, z_{N-1} = \theta_N$

$$\begin{aligned}
\tilde{\Phi}^{\underline{\beta}}_{\underline{\alpha}}(\underline{\theta}, \underline{z}) &\to \left(\Omega\, \tilde{C}^{\beta_m}(\underline{\theta}, z_m) \cdots \tilde{C}^{\beta_N}(\underline{\theta}, z_N) \tilde{C}^{\beta_{N-1}}(\underline{\theta}, \theta_N) \cdots \tilde{C}^{\beta_1}(\underline{\theta}, \theta_2)\right)_{\underline{\alpha}} \qquad \text{(B.8)}\\
&= \prod_{j=N}^{m} \tilde{b}(\theta_1 - z_j) \left(\tilde{S}_{1N} \dots \tilde{S}_{12}\right)^{\hat{\underline{\beta}},1}_{\hat{\underline{\alpha}}} \left(\Omega \tilde{C}^{\beta_m}(\check{\underline{\theta}}, z_m) \cdots \tilde{C}^{\beta_N}(\check{\underline{\theta}}, z_N)\right)_{\check{\underline{\alpha}}}.
\end{aligned}$$

It has been used that due to the $SU(N)$ ice rule only the amplitude $b(\cdot)$ (because all $\beta > 1$) contributes to the S-matrices $S(\theta_1 - z_j)$ $(j \ge N)$ and $a(\cdot)$ to the S-matrices $S(\theta_2 - z_j), \dots, S(\theta_N - z_j)$ $(j \ge N)$, $S(\theta_i - z_1), \dots, S(\theta_i - z_{N-2})$ $(i \ge N)$ after having applied Yang-Baxter relations. One observes that the product of S-matrices in (38) together with the one in (B.8) yields the total $N-1$ particle S-matrix

$$\tilde{S}^{N\dots 2}_{\hat{\underline{\beta}}}(\hat{\underline{z}}) \left(\tilde{S}_{1N} \dots \tilde{S}_{12}\right)^{\hat{\underline{\beta}},1}_{\hat{\underline{\alpha}}} = \tilde{S}^{N\dots 21}_{\hat{\underline{\alpha}}}(\hat{\underline{\theta}})$$

for which the residue formula (A.7) applies

$$\operatorname*{Res}_{\theta_{12}=i\eta}\dots\operatorname*{Res}_{\theta_{N-1N}=i\eta}\tilde S^{N\dots21}_{\alpha_1\dots\alpha_N}=(N-1)!(i\eta)^{N-1}\epsilon^{1\dots N}\epsilon_{\alpha_1\dots\alpha_N}\,.$$

We combine (B.8) with the function $L_{\underline{\beta}}(\underline{z})$ with the property (38) for $(l=1)$ and the scalar functions $\tilde h$ and p and after having performed the remaining z_j-integrations we obtain

$$R^{(1)}_{1\dots n}=c_0\prod_{i=N+1}^{n}\prod_{j=2}^{N}\tilde\phi(\theta_{ij})\,\varepsilon_{1\dots N}(\underline{\hat\theta})K_{N+1\dots n}(\underline{\check\theta})$$

$$c_0=c_1\nu(N-1)!(i\eta)^{N-1}(-1)^{m-N+1}\prod_{j=1}^{N-1}\tilde\phi(ji\eta)$$

We have used the equations (12), (17), the relation (iii') of (40) for the p-function, the relation (41) for the constant ν and the recursion relation (43) for the constants c_l with the solution (C.2).

The remaining contribution to (B.7) is due to R_2 and R_3

$$R^{(2)}_{1\dots n}+R^{(3)}_{1\dots n}=-c_0\prod_{i=N+1}^{n}\prod_{j=2}^{N}\tilde\phi(\theta_{ij})\,\varepsilon_{1\dots N}K_{N+1\dots n}(\underline{\check\theta})\dot\sigma_N S_{Nn}\dots S_{NN+1}$$

It is convenient to shift the particle with momentum θ_N to the right by applying S-matrices and write the claim as

$$\left(R^{(2)}_{1\dots n}+R^{(3)}_{1\dots n}\right)S_{N+1N}\dots S_{nN}+c_0\prod_{i=N+1}^{n}\prod_{j=2}^{N}\tilde\phi(\theta_{ij})\,\varepsilon_{1\dots N}K_{N+1\dots n}(\underline{\check\theta})\dot\sigma_N=0 \quad \text{(B.9)}$$

where the components of this co-vector are now denoted by $v_{\alpha_1\dots\alpha_{N-1}\alpha_{N+1}\dots\alpha_n\alpha_N}$. Note that because of (i) (see (30))

$$\begin{aligned}&K_{1\dots n}(\underline{\theta})S_{N+1N}\dots S_{nN}\\&=\prod_{i=N+1}^{n}a(\theta_{iN})K_{1\dots N-1N+1\dots nN}(\theta_1,\dots,\theta_{N-1},\theta_{N+1},\dots,\theta_n,\theta_N).\end{aligned}$$

Because the Bethe ansatz states are of highest weight (see [36]) it is sufficient to prove (B.9) only for $\alpha_N=1$. The contribution of $R^{(2)}_{1\dots n}$ is given by the $z_1,\dots,z_{N-1}$-integrations along the small circles around $z_1=\theta_1,\dots,z_{N-1}=\theta_{N-1}$ (see again figure 1). Now $\tilde S(\theta_1-z_1)_1,\dots,\tilde S(\theta_{N-1}-z_{N-1})$ yield permutation operators $\mathbf{P}$ and the co-vector part of this contribution for $\alpha_N=1$ is

$$\begin{aligned}&\left(\Omega\,\tilde C^{\beta_m}(\underline{\theta},z_m)\cdots\tilde C^{\beta_{N-1}}(\underline{\theta},\theta_{N-1})\cdots\tilde C^{\beta_1}(\underline{\theta},\theta_1)P_N(1)\right)_{\alpha_1\dots\alpha_{N-1}\alpha_{N+1}\dots\alpha_n\alpha_N}\\&\qquad=\delta^{\beta_1}_{\alpha_1}\cdots\delta^{\beta_{N-1}}_{\alpha_{N-1}}\left(\Omega\tilde C^{\beta_m}(\underline{\check\theta},z_m)\cdots\tilde C^{\beta_N}(\underline{\check\theta},z_N)\right)_{\underline{\check\alpha}}\delta^1_{\alpha_N}\end{aligned} \quad \text{(B.10)}$$

where $P_N(1)$ projects onto the components with $\alpha_N=1$. We have used the fact that because of the $SU(N)$ ice rule the amplitude $a(\cdot)$ only contributes to the S-matrices $S(\theta_1-z_j),S(\theta_2-z_j),S(\theta_N-z_j),S(\theta_i-z_1)$ after having applied Yang-Baxter relations. We use $\tilde\phi(\theta)=-\tilde b(\theta+2\pi i)\tilde\phi(\theta+2\pi i)$ to replace for $i=1,\dots,N-1$ and $\beta\neq1$

$$\tilde\phi(\theta_{Ni})=\tilde S^{11}_{11}(\theta_{Ni})\tilde\phi(\theta_{Ni})=-\tilde S^{\beta1}_{1\beta}(\theta_{Ni}+2\pi i)\tilde\phi(\theta_{Ni}+2\pi i)$$

therefore using again (A.7)

$$\operatorname*{Res}_{\theta_{N-1N}=i\eta}\cdots\operatorname*{Res}_{\theta_{12}=i\eta}\tilde\phi(\theta_{N1})\cdots\tilde\phi(\theta_{NN-1})\tilde S^{N\dots2}_{\alpha_1\dots\alpha_{N-1}}(\theta_1,\cdots,\theta_{N-1})$$
$$=-\tilde\phi(i\eta)\cdots\tilde\phi((N-1)i\eta)(N-1)!(i\eta)^{N-1}\epsilon_{\alpha_1\cdots\alpha_{N-1}1}\,.$$

We combine (B.10) with the function $L_{\underline\beta}(\underline z)$ with the property (38) (for $l=1$) and the scalar functions $\tilde h$ and p and after having performed the remaining z_j-integrations we obtain

$$R^{(2)}_{1\dots n}S_{N+1N}\dots S_{nN}P_N(1)=-c_0\dot\sigma\prod_{i=N+1}^{n}\prod_{j=2}^{N}\tilde\phi(\theta_{ij})\,\varepsilon_{1\dots N}K_{N+1\dots n}(\underline{\check\theta})P_N(1)\,. \tag{B.11}$$

Again we have used the equations (12), (17), the relation (iii') of (40) for the p-function, the relation (41) for the constant ν and the recursion relation (43) for the constants c_l with the solution (C.2).

The contribution of pinching (3) is given by the $z_1,\dots,z_{N-1}$-integrations along the small circles around $z_1=\theta_2-i\eta,\dots,\ z_{N-2}=\theta_{N-1}-i\eta,\ z_{N-1}=\theta_N-i\eta$, (see again figure 1). In $\tilde C^{\beta_{N-1}}_{\underline{\tilde\alpha}}(\underline\theta,z_{N-1})$ the S-matrix $\tilde S^{\beta'_{N-1}\alpha''_N}_{\alpha'_N 1}(\theta_N-z_{N-1})$ yields $\Gamma^{\beta'_{N-1}\alpha''_N}_{(\rho\sigma)}\Gamma^{(\rho\sigma)}_{\alpha'_N1}$ by applying $Res_{z_{N-1}=\theta_N-i\eta}$. For $\alpha_N=1$ this vanishes because $\Gamma^{(\rho\sigma)}_{\alpha\beta}$ is antisymmetric with respect to α,β the $SU(N)$ ice rule implies that also $\alpha'_N=1$. Therefore (B.9) is proved for $\alpha_N=1$ and because the left hand side belongs to a highest weight state also in general. ■

Appendix C. Proof of lemma 2

Proof. We set $n_l=m,\ n_{l+1}=k$. For the higher level functions $L^{(l)}_{\underline\beta}(\underline z)$ one may verify the equations (i)$^{(l)}$ and (ii)$^{(l)}$ quite analogously to the corresponding ones for the main theorem. We prove (iii)$^{(l)}$ by induction and assume:

$$L^{(l+1)}_{\underline\gamma}(\underline u)\approx c_{l+1}\tilde S^{N\dots l+2}_{\underline{\hat\gamma}}(\underline{\hat u})\prod_{i=N-l}^{k}\prod_{j=2}^{N-l-1}\tilde\phi(u_{ij})L^{(l+1)}_{\underline{\check\gamma}}(\underline{\check u}) \tag{C.1}$$

for $u_{12},\dots,u_{N-l-2,N-2}\to i\eta$. In the integral representation (34) of $L^{(l)}_{\underline\beta}(\underline z)$ there are pinchings at $u_1=z_2,\dots,u_{N-l-1}=z_{N-l}$ if $z_{12},\dots,z_{N-l-1,N-l}\to i\eta$. Therefore in $\tilde\Psi^{(l)}_{\underline\beta}(\underline z,\underline u)$ the S-matrices $\tilde S(z_2-u_1),\dots,\tilde S(z_{N-l}-u_{N-l-1})$ yield the permutation operator $\mathbf P$ and we have to consider

$$\tilde\Phi^{(l)\underline\gamma}_{\underline\beta}(\underline z,\underline u)=\Big(\Omega^{(l)}\,\tilde C^{(l)\gamma_k}(\underline z,u_k)\cdots\tilde C^{(l)\gamma_{N-l-1}}(\underline z,z_{N-l})\cdots\tilde C^{(l)\gamma_1}(\underline z,z_2)\Big)_{\underline\beta}$$
$$=\Big(\tilde S_{1N-l}(z_{1N-l})\dots\tilde S_{12}(z_{12})\Big)^{\gamma_1\dots\gamma_{N-l-1}l+1}_{\beta_1\dots\beta_{N-l}}\prod_{j=N-l}^{k}\tilde b(z_1-u_j)\tilde\Phi^{(l)\underline{\check\gamma}}_{\underline{\check\beta}}(\underline{\check z},\underline{\check u})$$

where $\underline{\check\beta}=(\beta_{N-l+1},\dots,\beta_m),\ \underline{\check z}=(z_{N-l+1},\dots,z_m),\ \underline{\check u}=(u_{N-l},\dots,u_k)$. We may write for $\underline{\hat u}=(u_1,\dots,u_{N-l-1})=(z_2,\dots,z_{N-l})$

$$\tilde S^{N\dots l+2}_{\underline{\hat\gamma}}(\underline{\hat u})\Big(\tilde S_{1N-l}(z_{1N-l})\dots\tilde S_{12}(z_{12})\Big)^{\gamma_1\dots\gamma_{N-l-1}l+1}_{\beta_1\dots\beta_{N-l}}=\tilde S^{N\dots l+1}_{\underline{\hat\beta}}(\underline{\hat z})$$

with the notation $\underline{\hat{z}} = (z_1, \dots, z_{N-l})$. Therefore using the assumption (C.1) we obtain when $z_{12}, \dots, z_{N-l-1,N-l} \to i\eta$

$$L^{(l)}_{\underline{\beta}}(\underline{z}) \approx \frac{1}{\check{k}!} \oint_{z_2} \frac{du_1}{R} \cdots \oint_{z_{N-l}} \frac{du_{N-l-1}}{R} \int_{\mathcal{C}_{\underline{z}}} \frac{du_{N-l}}{R} \cdots \int_{\mathcal{C}_{\underline{z}}} \frac{du_k}{R}\, \tilde{h}(\underline{z}, \underline{u})$$
$$\times L^{(l+1)}_{\underline{\gamma}}(\underline{u})\, \tilde{\Phi}^{(l)\underline{\gamma}}_{\underline{\beta}}(\underline{z}, \underline{u}) = c_l \tilde{S}^{N\dots l+1}_{\underline{\hat{\beta}}}(\underline{\hat{z}}) \prod_{i=N-l+1}^{m} \prod_{j=2}^{N-l} \tilde{\phi}(z_{ij}) L^{(l)}_{\underline{\check{\beta}}}(\underline{\check{z}})$$

where $\check{k} = k - N + l + 1$. The following formulae have been used:

$$\tilde{b}(z_1 - u_j) \prod_{i=3}^{N-l} \tilde{\phi}(u_j - z_i) \prod_{i=1}^{N-l} \tilde{\phi}(z_i - u_j) \prod_{i=2}^{N-l} \tau(z_i - u_j) = -1\,,$$

the relation $(\mathrm{iii}')^{(l)}$ of (42) for the p-function and the recursion relation

$$c_l = c_{l+1} (-1)^{n_{l+1} - N + l + 1} \prod_{j=1}^{N-l-1} \tilde{\phi}(ji\eta).$$

The solution of this recursion relation with $c_{N-1} = 1$ is

$$c_1 = \tilde{\phi}^{N-2}(i\eta) \tilde{\phi}^{N-3}(2i\eta) \cdots \tilde{\phi}((N-2)i\eta) \prod_{l=1}^{N-2} (-1)^{n_{l+1} - N + l + 1}. \quad \text{(C.2)}$$

■

References

[1] D. J. Gross and Frank Wilczek. Ultraviolet behaviour of non-abelian gauge theories. *Phys. Rev. Lett.*, 30:1343–1346, 1973.

[2] H. David Politzer. Reliable perturbative results for strong interactions? *Phys. Rev. Lett.*, 30:1346–1349, 1973.

[3] Niklas Beisert, Gabriele Ferretti, Rainer Heise, and Konstantin Zarembo. One-loop QCD spin chain and its spectrum. *Nucl. Phys.*, B717:137–189, 2005.

[4] H. Bethe. On the theory of metals. 1. eigenvalues and eigenfunctions for the linear atomic chain. *Z. Phys.*, 71:205–226, 1931.

[5] C. N. Yang. Some exact results for the many body problems in one dimension with repulsive delta function interaction. *Phys. Rev. Lett.*, 19:1312–1314, 1967.

[6] Bill Sutherland. Model for a multicomponent quantum system. *Phys. Rev. B*, 12(9):3795–3805, Nov 1975.

[7] L. D. Faddeev, E. K. Sklyanin, and L. A. Takhtajan. The quantum inverse problem method. 1. *Theor. Math. Phys.*, 40:688–706, 1980.

[8] Hratchya M. Babujian. Correlation function in wznw model as a bethe wave function for the gaudin magnetics. *In 'Gosen 1990, Proceedings, Theory of elementary particles' 12-23. (see high energy physics index 29 (1991) No. 12257)*, pages 12–23, 1990.

[9] I. B. Frenkel and N. Yu. Reshetikhin. Quantum affine algebras and holonomic difference equations. *Commun. Math. Phys.*, 146:1–60, 1992.

[10] V. Schechtman and A. Varchenko. Arrangements of hyerplanes and lie algebra homology. *Invent. Math.*, 106:139–194, 1991.

[11] M. Karowski and P. Weisz. Exact form factors in (1+1)-dimensional field theoretic models with soliton behaviour. *Nucl. Phys.*, B139:455–476, 1978.

[12] F.A. Smirnov. Form factors in completely integrable models of quantum field theory. *Adv. Series in Math. Phys. **14**, World Scientific*, 1992.

[13] H. M. Babujian, A. Fring, M. Karowski, and A. Zapletal. Exact form factors in integrable quantum field theories: The sine-gordon model. *Nucl. Phys.*, B538:535–586, 1999.

[14] Fabian H. L. Essler and Robert M. Konik. Applications of massive integrable quantum field theories to problems in condensed matter physics. *In 'Shifman, M. (ed.) et al.: From fields to strings'*, 1:684–830, 2004.

[15] A. M. Tsvelik. Quantum field theory in condensed matter physics. *Cambridge, UK: Univ. Pr.*, page 332, 1995.
[16] A. O. Gogolin, A. A. Nersesyan, and A. M. Tsvelik. Bosonization in strongly correlated systems. *Cambridge, UK: Univ. Pr.*, 1999.
[17] J. Links, H.Q. Zhou, R. McKenzie, and M. Gould. Algebraic bethe ansatz method for the exact calculation of energy spectra and form factors: applications to models of bose-einstein condensates and metallic nanograins. *J. Phys.*, A36:R63–R104, 2003.
[18] Davide Controzzi and Fabian H. L. Essler. Dynamical density correlation function of 1d mott insulators in a magnetic field. *Phys. Rev.*, B66:165112, 2002.
[19] Fabian H. L. Essler and Robert M. Konik. Dynamical spin response of doped two-leg hubbard-like ladders. *cond-mat/0607783*, 2004.
[20] J. Links and H.Q. Zhou. Exact form factors for the josephson tunneling current and relative particle number fluctuations in a model of two coupled bose-einstein condensates. *Lett. Math. Phys*, 60:275–282, 2002.
[21] J. L. Cardy, O. A. Castro-Alvaredo, and B. Doyon. Form factors of branch-point twist fields in quantum integrable models and entanglement entropy. *arXiv:0706.3384 [hep-th]*, 2007.
[22] H. Babujian and M. Karowski. Exact form factors for the scaling z(n)-ising and the affine a(n-1) toda quantum field theories. *Phys. Lett.*, B575:144–150, 2003.
[23] H. Babujian, A. Foerster, and M. Karowski. Exact form factors in integrable quantum field theories: The scaling z(n)-ising model. *Nucl. Phys.*, B736:169–198, 2006.
[24] Atsushi Nakayashiki and Yoshihiro Takeyama. On form factors of su(2) invariant thirring model. *math-ph/0105040*, 2001.
[25] Yoshihiro Takeyama. Form factors of su(n) invariant thirring model. *Publ. Res. Inst. Math. Sci. Kyoto*, 39:59–116, 2003.
[26] E. Mukhin, V. Tarasov, and A. Varchenko. Bethe eigenvectors of higher transfer matrices. *J. Stat. Mech.*, 06:P08002, 2006.
[27] S Pakuliak. Weight functions and nested bethe ansatz. *Ann. Hernri Poincare*, 7:1541–1554, 2007.
[28] M.J. Bhaseen and A.M. Tsvelik. SU(N) evolution of a frustrated spin ladder. *Physical Review B*, 68(9):94405, 2003.
[29] B. Berg, M. Karowski, V. Kurak, and P. Weisz. Factorized u(n) symmetric s matrices in two-dimensions. *Nucl. Phys.*, B134:125–132, 1978.
[30] V. Kurak and J. A. Swieca. Anti-particles as bound states of particles in the factorized s-matrix framework. *Phys. Lett.*, B82:289–291, 1979.
[31] R. Koberle, V. Kurak, and J. A. Swieca. Scattering theory and 1/n expansion in the chiral gross- neveu model. *Phys. Rev.*, D20:897–902, 1979.
[32] M. Karowski. On the bound state problem in (1+1) dimensional field theories. *Nucl. Phys.*, B153:244–252, 1979.
[33] H. Babujian and M. Karowski. Exact form factors in integrable quantum field theories: The sine-gordon model. ii. *Nucl. Phys.*, B620:407–455, 2002.
[34] H. Babujian, A. Foerster, and M. Karowski. Exact form factors in integrable quantum field theories: The su(n)-gross-neveu model. *to be published*, 2008.
[35] V. P. Yurov and Al. B. Zamolodchikov. Truncated fermionic space approach to the critical 2-d ising model with magnetic field. *Int. J. Mod. Phys.*, A6:4557–4578, 1991.
[36] Hratchia M. Babujian, Angela Foerster, and Michael Karowski. The nested su(n) off-shell bethe ansatz and exact form factors. *hep-th/0611012*, 2006.
[37] L. A. Takhtajan and L. D. Faddeev. The quantum method of the inverse problem and the heisenberg xyz model. *Russ. Math. Surveys*, 34:11–68, 1979.
[38] H. Babujian and M. Karowski. Sine-gordon form factors and quantum field equations. *J. Phys.*, A35:9081–9104, 2002.
[39] H. Babujian, M. Karowski, and A. Zapletal. Matrix difference equations and a nested bethe ansatz. *J. Phys.*, A30:6425–6450, 1997.
[40] David J. Gross and Andre Neveu. Dynamical symmetry breaking in asymptotically free field theories. *Phys. Rev.*, D10:3235, 1974.
[41] H. Babujian, M. Karowski, and A. Zapletal. U(n) matrix difference equations and a nested bethe ansatz. *hep-th/9611006*, 1996.
[42] R. Köberle and J. A. Swieca. Factorizable z(n) models. *Phys. Lett.*, B86:209–210, 1979.
[43] H. Babujian, M. Karowski, and A. Zapletal. Su(n) matrix difference equations and a nested bethe ansatz. *hep-th/9611005*, 1996.