

# Probing strong correlations with light scattering: Example of the quantum Ising model

H. M. Babujian,<sup>1</sup> M. Karowski,<sup>2</sup> and A. M. Tselik<sup>3</sup>

<sup>1</sup>*Yerevan Physics Institute, Alikhanian Brothers 2, Yerevan 375036, Armenia;*

*International Institute of Physics, Universidade Federal do Rio Grande do Norte (UFRN), 59078-400 Natal-RN, Brazil;*  
*and Simons Center for Geometry and Physics, Stony Brook University, Stony Brook, New York 11794, USA*

<sup>2</sup>*Institut für Theoretische Physik, Freie Universität Berlin, Arnimallee 14, 14195 Berlin, Germany*

<sup>3</sup>*Condensed Matter Physics and Materials Science Division, Brookhaven National Laboratory, Upton, New York 11973-5000, USA*

(Received 27 May 2016; revised manuscript received 17 October 2016; published 31 October 2016)

In this paper we calculate the nonlinear susceptibility and the resonant Raman cross section for the paramagnetic phase of the ferromagnetic quantum Ising model in one dimension. In this region the spectrum of the Ising model has a gap  $m$ . The Raman cross section has a strong singularity when the energy of the outgoing photon is at the spectral gap  $\omega_f \approx m$  and a square root threshold when the frequency difference between the incident and outgoing photons  $\omega_i - \omega_f \approx 2m$ . The latter feature reflects the fermionic nature of the Ising model excitations.

DOI: [10.1103/PhysRevB.94.155156](https://doi.org/10.1103/PhysRevB.94.155156)

## I. INTRODUCTION

When photons enter into a strongly interacting medium one expects nonlinear phenomena such as frequency mixing and inelastic light scattering [1,2]. These effects are interesting by themselves, but they can also serve as experimental tools to extract otherwise inaccessible information about strongly correlated dynamics since all these tools probe multipoint dynamical correlation functions which carry much richer information than a simple linear response.

In this paper we will present calculations of three- and four-point dynamical correlation functions for the ferromagnetic quantum Ising (FQI) model in one spatial dimension (1D) and relate them to two spectroscopic probes: the nonlinear susceptibility and the inelastic scattering cross section of light (Raman scattering). We have chosen the 1D FQI model for three reasons. First, this is a strongly correlated model whose applicability to real materials has been firmly established (see the discussion at the end of the paper). Secondly, this model admits a resonance regime of where nonlinear effects are strongly enhanced. The effectiveness of light probes is somewhat restricted by the fact that the photons carry a very small momentum, but in most strongly correlated systems the strongest correlations occur at finite wave vectors of order of the size of the Brillouin zone. The 1D FQI model represents an exception from this rule. The third factor is a comparative simplicity of the calculations which will allow us to concentrate on the crucial points avoiding cumbersome technical details. We consider FQI as the first integrable model among many where multipoint correlation functions can be calculated and their current calculations present a skeleton of the general scheme for all such models.

The quantum Ising model is described by the Hamiltonian

$$H = \sum_n (-J\sigma_n^z\sigma_{n+1}^z + h\sigma_n^x), \quad (1)$$

where  $\sigma^a$  are the Pauli matrices. This model is one of the best studied models of strongly correlated physics (see [3–5] for a review). Its exact solution was obtained as early as 1928 by Jordan-Wigner transformation which expresses the spin operators in terms of fermionic creation and annihilation operators. For the reasons explained above we are interested

in the case  $J > 0$  (FQI model). The best condensed matter realization known to date is found in columbite  $\text{CoNb}_2\text{O}_6$  [6–8]. We will discuss it in the Conclusions.

The Ising model (1) may describe not only spins, but any coupled two level systems. If these are spins, then operators  $\sigma_n^z$  directly couple to external magnetic field:  $\mu_B B_n^z \sigma_n^z$ . Alternatively the states of two level system may correspond to positions of electric charges in a double-well potential. Then  $\sigma^a$  would be the dipole moment operators; the first term in (1) is the dipole-dipole interaction and the transverse field describes the quantum tunneling between the wells. The interaction of dipole moments with the electric field is given by  $pE_n^z \sigma_n^z$  with  $p$  being the dipole moment magnitude.

When the dominant interaction is ferromagnetic  $J > 0$ , the strongest fluctuations take place at zero wave vectors which guarantees a direct coupling to the electromagnetic field creating optimal resonance conditions. The Ising model (1) has two phases depending on the sign of  $m = h - J$ . The resonance occurs in the paramagnetic phase  $m > 0$  when the ground state average of the order parameter  $\langle \sigma^z \rangle = 0$ . In that case the electromagnetic field has a nonzero matrix element between the ground state and single magnon state.

Raman light scattering is a powerful experimental technique frequently used in condensed matter physics. The measured quantity is the inelastic scattering cross section of photons  $R(\mathbf{q}, \Omega)$  which contains information about the excitations of condensed matter systems with which the photons interact. The theory of Raman scattering was formulated in the 1920s [9,10] when the formulas for  $R(\mathbf{q}, \Omega)$  were derived (see also [11]). A radical simplification of these formulas was suggested in [12] where the resonant part of the Raman cross section was expressed as a particular limit of the four-point correlation function of the current operators. This simplification allows one to apply to the problem various techniques of quantum field theory such as Feynman diagram expansion and also simplifies the application of nonperturbative techniques.

Another technique to be discussed is a nonlinear response directly related to the three-point correlation function. The related themes are two-dimensional spectroscopy and spectroscopy with entangled photons [13,14]. A relation of third-order nonlinear optical properties to magnetic interactions was demonstrated in [15].

The Jordan-Wigner transformation transforms FQI into a model of noninteracting massive Majorana fermions. In the scaling limit  $m \ll J$  their dispersion becomes relativistic  $\epsilon(k) = \sqrt{v^2 k^2 + m^2}$ ,  $v \sim J$ . In what follows we will set  $v = 1$  to restore  $J$  in the final expressions.

The fact that the excitations of the Ising model do not interact does not make the model trivial. Indeed, since  $\sigma^z$  operators are very nonlocal in terms of the fermions, the electromagnetic field has matrix elements between states with different number of fermionic excitations. Such situation is typical for strongly interacting systems and experimental probes of multipoint correlators are highly suitable to reveal this nonlocality. In the paramagnetic phase of FQI the inelastic processes involve matrix elements with odd number of the Ising fermions with the leading low energy processes being transitions from single- to two-fermion states. The fermionic nature of the excitations is reflected in the fact that the cross section vanishes at the threshold:  $R(\Omega) \sim (\Omega - 2m)^{1/2}$  [see Eqs. (8) and (9) below.

## II. THE OBSERVABLES

### A. The nonlinear susceptibility

The nonlinear susceptibility is the third derivative of the action with respect to the dynamical magnetic field:

$$\chi_{zzx}^{(3)} = \langle \hat{T} \sigma^z(t_1) \sigma^z(t_2) \sigma^x(t_3) \rangle_{\text{connected}}, \quad (2)$$

where  $\sigma = \sum_n \sigma_n^z$  (we assume that electromagnetic radiation carries no momentum). In the Ising model such response exists only when the polarization of photons is such that the magnetic field has both  $z$  and  $x$  components so that the coupling to the magnetic field is described as

$$V = \mu_B \sum_n (B^z \sigma_n^z + B^x \sigma_n^x). \quad (3)$$

The nonlinear susceptibility describes the effects of frequency mixing. As follows from (2), the only nonzero third order response includes two magnetic fields with frequencies  $\omega_z, \omega_x - \omega_z$  along the  $z$  direction and one field with frequency  $-\omega_x$  along the  $x$  direction. Our result where we take into account only two magnon production processes, is

$$\begin{aligned} \chi_{zzx}^{(3)}(\omega_z, -\omega_z + \omega_x, -\omega_x) \\ = C^2 \mu_B^3 (mJ^3)^{-1/4} \left\{ \frac{(\omega_x - 2m)h(\omega_x/2m + i\delta)}{(\omega_z - m)(\omega_x - \omega_z - m)} \right. \\ \left. + (\omega_{x,z} \rightarrow -\omega_{x,z}) \right. \\ \left. + 16m\pi \frac{(\omega_x - \omega_z)\omega_z - m^2}{(\omega_z^2 - m^2)[(\omega_x - \omega_z)^2 - m^2]} \right\}, \quad (4) \end{aligned}$$

where  $C \sim 1$  is a numerical constant and (see Appendix C)

$$h(x) = \int_{-\infty}^{\infty} \left( 1 + \frac{1}{\cosh \theta} \right)^2 \frac{1}{\cosh \theta - x} d\theta.$$

### B. Relation between the Raman cross section and the correlation functions

As was stated above we assume the following spin-photon interaction  $V = \sum_n p E_n \sigma_n^z$ . Then according to Eq. (2.21)

from [12] the cross section for the light beam polarized along the  $z$  axis is given by the following expression:

$$R(\omega_i, \omega_f) = 2\pi \mu_B^4 [(h\omega_i)(h\omega_f)] \frac{\chi_R(\omega_i, \omega_f)}{1 - \exp[-\beta(\omega_i - \omega_f)]}, \quad (5)$$

where  $\beta = 1/T$ , and  $\omega_i$  and  $\omega_f$  are frequencies of the incident and the scattered light. We will consider the  $T = 0$  limit. Then the function  $\chi$  is expressed as (see Eqs. (2.30) and (2.31) from [12])

$$\begin{aligned} \chi_R(\omega_i, \omega_f) = \frac{1}{2\pi i} \lim_{\delta_1 > \delta_2 \rightarrow 0} \{ \tilde{\Xi}(-\omega_i - i\delta_1, \omega_f + i\delta_2, -\omega_f \\ + i\delta_2, \omega_i - i\delta_1) - \tilde{\Xi}(-\omega_i - i\delta_2, \omega_f \\ + i\delta_1, -\omega_f + i\delta_1, \omega_i - i\delta_2) \}, \quad (6) \end{aligned}$$

where  $\tilde{\Xi}$  is the Fourier transform of the four-point time ordered correlation function:

$$\langle \hat{T} \sigma(t_1) \sigma(t_2) \sigma(t_3) \sigma(t_4) \rangle_{\text{connected}}. \quad (7)$$

Below we will derive the expression for (7) in the paramagnetic phase of model (1) at  $T = 0$  in the limit  $m = h - J \ll J$  and will use the result to calculate the Raman cross section (5). For  $\omega_i > \omega_f > 0$  we obtain the result by substituting Eq. (20) into (6) (see Appendix B)

$$\begin{aligned} \chi_R(\omega_i, \omega_f) \sim (m/J)^{1/2} \frac{J^3}{m^4} \left\{ \frac{G\left[\frac{1}{2m}(\omega_i + \omega_f)\right]}{(\omega_f - m)^2(\omega_i - m)^2} \right. \\ \left. + \frac{G\left[\frac{1}{2m}(\omega_i - \omega_f)\right]}{(\omega_f + m)^2(\omega_i - m)^2} \right\}, \quad (8) \end{aligned}$$

where

$$G(x) = \Theta(x - 1) \frac{(x - 1)^{1/2}(x + 1)^{5/2}}{x^3}. \quad (9)$$

The calculation takes into account only two-particle intermediate states and hence is valid in the range of frequencies  $|\omega_i - \omega_f| < 4m$  when the processes with emission of more than two particles do not contribute to the inelastic cross section. The threshold for the inelastic scattering is at  $\omega_i \pm \omega_f = 2m$  corresponding to the emission of two fermionic excitations.

## III. GREEN'S FUNCTIONS

Below we will do our calculations in the most general form valid for all integrable models and at the end apply the results to the Ising model. We will concentrate on the most difficult case of the four-point function; the calculations of the three-point one are comparatively straightforward.

Let  $\phi$  be a scalar bosonic field (for the Ising model it is the scaling limit of the  $\sigma^z$ ). Its Green's functions are defined as time ordered  $n$ -point functions

$$\begin{aligned} \tau(\underline{x}) &= \langle 0 | T \phi(x_1) \cdots \phi(x_n) | 0 \rangle \\ &= \sum_{\text{perm}(\underline{x})} \Theta_{1 \dots n}(\underline{t}) w(\underline{x}). \end{aligned}$$

Here  $w(\underline{x}) = \langle 0 | \phi(x_1) \cdots \phi(x_n) | 0 \rangle$  is the Wightman function and  $\Theta_{1 \dots n}(\underline{t}) = \Theta(t_{12}) \Theta(t_{23}) \cdots \Theta(t_{n-1, n})$ . In momentum

space

$$\begin{aligned}\tilde{\tau}(\underline{k}) &= \int \underline{d^2x} e^{i x_i k_i} \tau(\underline{x}) \\ &= \sum_{\text{perm}(k)} \int \underline{d^2x} e^{i x_i k_i} \Theta_{1\dots n} w(\underline{x}).\end{aligned}$$

The connected Green's functions are given by

$$\tilde{\tau}(\underline{k}) = \sum_{\underline{k}_1 \cup \dots \cup \underline{k}_m = \underline{k}} \tilde{\tau}_c(\underline{k}_1) \cdots \tilde{\tau}_c(\underline{k}_m). \quad (10)$$

For convenience we split off the energy momentum  $\delta$  function and define  $\tilde{\Pi}(\underline{k})$  by

$$\tilde{\tau}_c(\underline{k}) = (2\pi)^2 \delta^{(2)}(\sum k_i) \tilde{\Xi}(\underline{k}). \quad (11)$$

*S-matrix and form factors.* For integrable quantum field theories the  $n$ -particle  $S$  matrix factorizes into  $n(n-1)/2$  two-particle ones

$$S^{(n)}(\theta_1, \dots, \theta_n) = \prod_{i < j} S(\theta_{ij}),$$

where the product on the right-hand side has to be taken in a specific order (see, e.g., [16]). The numbers  $\theta_{ij}$  are the rapidity differences  $\theta_{ij} = \theta_i - \theta_j$ , which are related to the momenta of the particles by  $p_i = m(\cosh \theta_i, \sinh \theta_i)$ . The form factors of a bosonic field are defined as the matrix elements

$$F(\underline{\theta}) = \langle 0 | \varphi(0) | \theta_1, \dots, \theta_n \rangle \quad (12)$$

(for the paramagnetic phase of the Ising model they are nonzero for  $n = \text{odd}$ ). They satisfy the form factor equations (i)–(v) (see, e.g., [17]). We use the normalization  $\langle 0 | \varphi(0) | \theta \rangle = 1$ . As a generalization we write

$$F(\underline{\theta}'; \underline{\theta}) = \langle \theta'_1, \dots, \theta'_n | \varphi(0) | \theta_1, \dots, \theta_n \rangle, \quad (13)$$

which is related to (12) by crossing. In particular (see Appendix A),

$$F(\theta_1; \theta_2, \theta_3) = F(\theta_1, \theta_2 - i\pi_-, \theta_3 - i\pi_+) + \delta_{\theta_{12}} + \delta_{\theta_{13}}, \quad (14)$$

$$F(\theta_2, \theta_3; \theta_4) = F(\theta_3 + i\pi_+, \theta_2 + i\pi_-, \theta_4) + \delta_{\theta_{24}} + \delta_{\theta_{34}} \quad (15)$$

with  $i\pi_{\pm} = i\pi \pm i\epsilon$  and  $\delta_{\theta_{12}} = 4\pi\delta(\theta_1 - \theta_2)$ .

### A. The Green's functions in the low particle approximation

Calculations of the Green's functions of massive theories simplify when we restrict our interest to their imaginary parts, as in the case of Raman scattering (5). In that case for any given energy only a limited number of matrix elements contribute to the calculations, namely, those ones which correspond to emissions of particles whose energy does not exceed the threshold.

Below we will derive expressions for the four-point Green's functions in the two-particle approximation.

The two-point Wightman function in one-particle intermediate states approximation is (with the short notation

$$\int_{\theta} = \frac{1}{4\pi} \int d\theta)$$

$$\begin{aligned}w^1(x_1 - x_2) &= \int_{\theta} \langle 0 | \varphi(x_1) | \theta \rangle \langle \theta | \varphi(x_2) | 0 \rangle \\ &= \int \frac{dp}{2\pi 2\omega} e^{-i(x_1 - x_2)p} = i\Delta_+(x_1 - x_2).\end{aligned}$$

The four-point Wightman function in 1-0-1 intermediate particle approximation is

$$\begin{aligned}w^{101}(\underline{x}) &= \int_{\theta_1} \langle 0 | \varphi(x_1) | \theta_1 \rangle \langle \theta_1 | \varphi(x_2) | 0 \rangle \\ &\quad \times \int_{\theta_4} \langle 0 | \varphi(x_3) | \theta_4 \rangle \langle \theta_4 | \varphi(x_4) | 0 \rangle \\ &= w^1(x_1 - x_2) w^1(x_3 - x_4).\end{aligned} \quad (16)$$

The four-point Wightman function in 1-2-1 intermediate particle approximation is [with  $\int_{\underline{\theta}} = \int_{\theta_1} \cdots \int_{\theta_4}$  and  $\mathbf{x}p = x_1 p_1 + x_2(p_2 + p_3 - p_1) + x_3(p_4 - p_2 - p_3) - x_4 p_4$ ]

$$\begin{aligned}w^{121}(\underline{x}) &= \frac{1}{2} \int_{\underline{\theta}} \langle 0 | \varphi(x_1) | \theta_1 \rangle \langle \theta_1 | \varphi(x_2) | \theta_2, \theta_3 \rangle \\ &\quad \times \langle \theta_3, \theta_2 | \varphi(x_3) | \theta_4 \rangle \langle \theta_4 | \varphi(x_4) | 0 \rangle \\ &= \frac{1}{2} \int_{\underline{\theta}} e^{-i\mathbf{x}p} F(\theta_1; \theta_2, \theta_3) F(\theta_2, \theta_3; \theta_4)\end{aligned}$$

with [see (14) and (15)]

$$\begin{aligned}\frac{1}{2} F(\theta_1; \theta_2, \theta_3) F(\theta_2, \theta_3; \theta_4) &= \frac{1}{2} (F(\theta_1, \theta_2 - i\pi_-, \theta_3 - i\pi_+) + \delta_{\theta_{12}} + \delta_{\theta_{13}}) \\ &\quad \times (F(\theta_3 + i\pi_+, \theta_2 + i\pi_-, \theta_4) + \delta_{\theta_{24}} + \delta_{\theta_{34}}) \\ &= I(\underline{\theta}) = I_1(\underline{\theta}) + I_2(\underline{\theta}) + I_3(\underline{\theta}).\end{aligned}$$

We have introduced (see Appendix A)

$$\begin{aligned}I_1(\underline{\theta}) &= \frac{1}{4} F(\theta_1, \theta_2 - i\pi_+, \theta_3 - i\pi_-) \\ &\quad \times F(\theta_3 + i\pi_+, \theta_2 + i\pi_-, \theta_4) \\ &\quad + \frac{1}{4} F(\theta_1, \theta_2 - i\pi_-, \theta_3 - i\pi_+) \\ &\quad \times F(\theta_3 + i\pi_-, \theta_2 + i\pi_+, \theta_4), \\ I_2(\underline{\theta}) &= \frac{1}{4} \{ \delta_{\theta_{12}} [1 + S(\theta_{23})] + \delta_{\theta_{13}} [1 + S(\theta_{23})] \} \\ &\quad \times F(\theta_3 + i\pi_+, \theta_2 + i\pi_-, \theta_4) \\ &\quad + \frac{1}{4} F(\theta_1, \theta_2 - i\pi_-, \theta_3 - i\pi_+) \\ &\quad \times \{ \delta_{\theta_{24}} [1 + S(\theta_{32})] + \delta_{\theta_{34}} [1 + S(\theta_{32})] \}, \\ I_3(\underline{\theta}) &= \frac{1}{2} (\delta_{\theta_{12}} + \delta_{\theta_{13}}) (\delta_{\theta_{24}} + \delta_{\theta_{34}}).\end{aligned} \quad (17)$$

From  $I_3$  we calculate

$$\begin{aligned}w_3^{121}(\underline{x}) &= w^1(x_1 - x_4) w^1(x_2 - x_3) \\ &\quad + w^1(x_1 - x_3) w^1(x_2 - x_4).\end{aligned} \quad (18)$$

Therefore neglecting contributions from higher particle intermediate states using (10) and (16) we obtain the connected four-point Green's function

$$\tilde{\tau}_c(\underline{k}) = \sum_{\text{perm}(k)} \int \underline{d^2x} \Theta_{1\dots n} e^{i x_i k_i} [w_1^{121}(\underline{x}) + w_2^{121}(\underline{x})], \quad (19)$$

where  $w_i^{121}(x)$  is given by the contribution from  $I_i(\underline{\theta})$  in (17). For  $k_i = (k_i^0, 0)$  we obtain from (11) (see Appendix B)

$$\tilde{\Xi}(k) = \frac{1}{32\pi m^6} \sum_{\text{perm}(k)} \frac{m}{m - k_1^0 - i\epsilon} \times \frac{m}{k_4^0 + m - i\epsilon} g\left(\frac{-1}{2m}(k_3^0 + k_4^0)\right) \quad (20)$$

with

$$g(x) = -2\pi \int_{\theta} \frac{1}{\omega/m} I(0, \theta, -\theta, 0) \frac{1}{\omega/m - x}, \quad (21)$$

where  $I = I_1 + I_2$  contribute. For integrable models typically  $S(0) = -1$ , then the contribution from  $I_2$  vanishes for  $\theta_i \rightarrow 0$ . With (6) and  $G(x) = (x - 1)^2 \text{Im } g(x)$  Eq. (8) follows. Next we consider a simple model, for which we calculate the function  $g(x)$  explicitly.

### B. The scaling limit of the Ising model

In the scaling limit this model may be described by an interacting Bose field  $\sigma_n^z = C m^{1/8} \phi(x)$ , where  $C$  is a numerical constant and  $m = h - J$ . The excitations are noninteracting Majorana fermions with the two-particle  $S$  matrix  $S(\theta) = -1$ . The field  $\sigma^x = (m/J)^{1/2} \epsilon(x) \sim \bar{\psi} \psi(x)$ , where  $\psi$  is a free Majorana spinor field. In [3,18,19] the form factor was proposed

$$F(\underline{\theta}) = \langle 0 | \sigma(0) | \theta_1, \dots, \theta_n \rangle = (2i)^{(n-1)/2} \prod_{i < j} \tanh \frac{1}{2} \theta_{ij}. \quad (22)$$

For  $k_i = (k_i^0, 0)$  in momentum space the contribution from  $I_2$  in (17) vanishes, because  $S(0) = -1$ . From (17) and (22) we obtain (see Appendix B)

$$I_1(0, \theta, -\theta, 0) = \tanh^2 \theta \coth^4 \frac{1}{2} (\theta - i\epsilon) + (\epsilon \rightarrow -\epsilon).$$

Substituting it into (21) and taking into account  $G(x) = (x - 1)^2 \text{Im } g(x)$  and the relation between  $\sigma^z$  and  $\phi$  we obtain (9).

From  $\epsilon(x) \sim \bar{\psi} \psi(x)$  one has for a free Majorana spinor field

$$\langle 0 | \epsilon(0) | \theta_1, \theta_2 \rangle = \sinh(\theta_{12}/2). \quad (23)$$

For low intermediate particle numbers this leads to (4) as above (see Appendix C).

## IV. CONCLUSIONS

We calculated the three- and the four-point correlation functions for the ferromagnetic quantum Ising model and discussed their relation to the observable quantities. In the paramagnetic phase of FQI the magnetic field is directly coupled to the spin operator which has matrix elements between states with odd and even numbers of Ising fermions. The fact that light can create an odd number of fermionic excitations is quite remarkable. It emphasizes an ambiguity between bosons and fermions existing in one dimension.

As we have said in Introduction, the best experimental realization of the FQI model known to date is found in columbite  $\text{CoNb}_2\text{O}_6$  [6–8]. Another possible candidate is  $\text{Sr}_3\text{CuIrO}_6$  [20]. Both these materials are quasi-1D insulators; the columbite displays a quantum critical point at  $B = 5.5$  T which is very well described by the theory of the Ising model [7]. Neutron scattering [6] and terahertz spectroscopy [8] also yield excellent agreement with the theoretical predictions. In view of these we suggest that a good test of our theory would be high field spectroscopic measurements at terahertz frequencies on  $\text{CoNb}_2\text{O}_6$ .

## ACKNOWLEDGMENTS

We are grateful to G. Blumberg, J. Misewich, and especially to N. P. Armitage for advising us on experimentally related matters, to S. Lukyanov who pointed out for the paper [21], and to A. B. Zamolodchikov for fruitful discussions. A.M.T. was supported by the U.S. Department of Energy (DOE), Division of Materials Science, under Contract No. DE-AC02-98CH10886. H.B. is grateful to Simons Center and Brookhaven National Laboratory for hospitality and support. H.B. is also supported by Armenian Grant No. 15T-1C308 and by ICTP OEA-AC-100 project. M.K. was supported by Fachbereich Physik, Freie Universität Berlin.

## APPENDIX A: CROSSING

The form factors (13) satisfy crossing relations (see, e.g., (31) in [22]), in particular

$$\begin{aligned} F(\theta_1; \theta_2, \theta_3) &= \langle \theta_1 | \varphi(0) | \theta_2, \theta_3 \rangle = F(\theta_1 + i\pi_-, \theta_2, \theta_3) + \delta_{\theta_{12}} + \delta_{\theta_{13}} S(\theta_{23}), \\ F(\theta_2, \theta_3; \theta_4) &= \langle \theta_3, \theta_2 | \varphi(0) | \theta_4 \rangle = F(\theta_3 + i\pi_-, \theta_2 + i\pi_-, \theta_4) + \delta_{\theta_{24}} + \delta_{\theta_{34}} S(\theta_{32}) \end{aligned} \quad (A1)$$

with  $i\pi_{\pm} = i\pi \pm i\epsilon$  and  $\delta_{\theta_{12}} = 4\pi \delta(\theta_1 - \theta_2)$ . Using the form factor equation (iii) and Lorentz invariance (see, e.g., [17])

$$\text{Res}_{\theta_{12}=i\pi} F(\theta_1, \theta_2, \theta_3) = 2i[\mathbf{1} - S(\theta_{23})], \quad F(\theta_1, \theta_2, \theta_3) = F(\theta_1 + \mu, \theta_2 + \mu, \theta_3 + \mu)$$

we can rewrite these equations as (14) and (15). And further one derives

$$\frac{1}{2} F(\theta_1, \theta_2 - i\pi_-, \theta_3 - i\pi_+) + \delta_{\theta_{12}} + \delta_{\theta_{13}} = \frac{1}{2} \{ F(\theta_1, \theta_2 - i\pi_+, \theta_3 - i\pi_-) + \delta_{\theta_{12}} [1 + S(\theta_{23})] + \delta_{\theta_{13}} [1 + S(\theta_{23})] \}, \quad (A2)$$

$$\frac{1}{2} F(\theta_3 + i\pi_+, \theta_2 + i\pi_-, \theta_4) + \delta_{\theta_{24}} + \delta_{\theta_{34}} = \frac{1}{2} \{ F(\theta_3 + i\pi_-, \theta_2 + i\pi_+, \theta_4) + \delta_{\theta_{24}} [1 + S(\theta_{32})] + \delta_{\theta_{34}} [1 + S(\theta_{32})] \}. \quad (A3)$$

Using Eqs. (14) and (15) and the identity  $(a + b + c)(d + e + f) = (\frac{1}{2}a + b + c)d + a(\frac{1}{2}d + e + f) + (b + c)(e + f)$  we derive

$$\begin{aligned} F(\theta_1; \theta_2, \theta_3)F(\theta_2, \theta_3; \theta_4) &= [F(\theta_1, \theta_2 - i\pi_-, \theta_3 - i\pi_+) + \delta_{\theta_{12}} + \delta_{\theta_{13}}][F(\theta_3 + i\pi_+, \theta_2 + i\pi_-, \theta_4) + \delta_{\theta_{24}} + \delta_{\theta_{34}}] \\ &= [\frac{1}{2}F(\theta_1, \theta_2 - i\pi_-, \theta_3 - i\pi_+) + \delta_{\theta_{12}} + \delta_{\theta_{13}}]F(\theta_3 + i\pi_+, \theta_2 + i\pi_-, \theta_4) \\ &\quad + F(\theta_1, \theta_2 - i\pi_-, \theta_3 - i\pi_+)[\frac{1}{2}F(\theta_3 + i\pi_+, \theta_2 + i\pi_-, \theta_4) + \delta_{\theta_{24}} + \delta_{\theta_{34}}] + (\delta_{\theta_1\theta_2} + \delta_{\theta_1\theta_3})(\delta_{\theta_4\theta_2} + \delta_{\theta_4\theta_3}), \end{aligned}$$

then (A2) and (A3) prove (17).

## APPENDIX B: PROOF OF (8)

### 1. Calculation of $\tau_c^{121}$

To derive (20) from (19) we calculate (for  $i = 1, 2$ )

$$\begin{aligned} &\int \underline{d^2x} \Theta_{1\dots n} e^{ix_i k_i} \int_p e^{-ix_1 p_1 - ix_2(p_2 + p_3 - p_1) - ix_3(p_4 - p_2 - p_3) + ix_4 p_4} I_i(\theta_1, \theta_2, \theta_3, \theta_4) \\ &= \int \underline{dx}^0 \Theta_{1\dots n} e^{ix_i^0 k_i^0} \int_\theta e^{-ix_1^0 \omega_1 - ix_2^0(\omega_2 + \omega_3 - \omega_1) - ix_3^0(\omega_4 - \omega_2 - \omega_3) + ix_4^0 \omega_4} \\ &\quad \times (2\pi)^4 \delta(p_1 - k_1^1) \delta(p_2 + p_3 - p_1 - k_2^1) \delta(p_4 - p_2 - p_3 - k_3^1) \delta(p_4 + k_4^1) I_i(\theta_1, \theta_2, \theta_3, \theta_4). \end{aligned}$$

For  $k_i = (k_i^0, 0)$  this is equal to

$$\begin{aligned} &2\pi \delta(k_1^1 + k_2^1 + k_3^1 + k_4^1) \int \underline{dx}^0 \Theta_{1\dots n} e^{ix_i^0 k_i^0} \int_\theta e^{-ix_1^0 m - ix_2^0(\omega_2 + \omega_3 - m) - ix_3^0(m - \omega_2 - \omega_3) + ix_4^0 m} (2\pi)^3 \delta(p_1) \delta(p_2 + p_3) \delta(p_4) I_i(0, \theta_2, -\theta_2, 0) \\ &= 2\pi \delta(k_1^1 + k_2^1 + k_3^1 + k_4^1) \frac{1}{(2m)^2} \int_\theta \frac{1}{2\omega} I_i(0, \theta, -\theta, 0) \int_{-\infty}^{\infty} dx_1^0 \int_{-\infty}^0 dx_2^0 \int_{-\infty}^0 dx_3^0 \int_{-\infty}^0 dx_4^0 \\ &\quad \times e^{ix_1^0(k_1^0 - m) + i(x_2^0 + x_1^0)[k_2^0 - (2\omega - m)] + i(x_3^0 + x_2^0 + x_1^0)[k_3^0 - (m - 2\omega)] + i(x_4^0 + x_3^0 + x_2^0 + x_1^0)(k_4^0 + m)} \end{aligned}$$

and

$$\begin{aligned} &\int_{-\infty}^{\infty} dx_1^0 \int_{-\infty}^0 dx_2^0 \int_{-\infty}^0 dx_3^0 \int_{-\infty}^0 dx_4^0 e^{ix_1^0(k_1^0 - m) + i(x_2^0 + x_1^0)[k_2^0 - (2\omega - m)] + i(x_3^0 + x_2^0 + x_1^0)[k_3^0 - (m - 2\omega)] + i(x_4^0 + x_3^0 + x_2^0 + x_1^0)(k_4^0 + m)} \\ &= 2\pi \delta(k_1^0 + k_2^0 + k_3^0 + k_4^0) \frac{-i}{k_2^0 + k_3^0 + k_4^0 + m - i\epsilon} \frac{-i}{k_3^0 + k_4^0 + 2\omega - i\epsilon} \frac{-i}{k_4^0 + m - i\epsilon} \end{aligned}$$

proves (20) and (21). For integrable models typically  $S(0) = -1$ , then the contribution from  $I_2$  vanishes for  $\theta_i \rightarrow 0$  [for the scaling Ising model we have  $S(\theta) \equiv -1$ ].

### 2. The function $g(x)$ for the scaling Ising model

From (17) and (22) we obtain (up to const)

$$\begin{aligned} I_1(0, \theta, -\theta, 0) &= \frac{1}{4} F(0, \theta - i\pi_+, -\theta - i\pi_-) F(-\theta + i\pi_+, \theta + i\pi_-, 0) + (\epsilon \rightarrow -\epsilon) \\ &= \{ [\tanh \frac{1}{2}(-\theta + i\pi + i\epsilon)] [\tanh \frac{1}{2}(\theta + i\pi - i\epsilon)] \tanh \frac{1}{2}(2\theta) \} \\ &\quad \times \{ [\tanh \frac{1}{2}(-2\theta)] [\tanh \frac{1}{2}(-\theta + i\pi + i\epsilon)] \tanh \frac{1}{2}(\theta + i\pi - i\epsilon) \} + (\epsilon \rightarrow -\epsilon) \\ &= \tanh^2 \theta \coth^4 \frac{1}{2}(\theta - i\epsilon) + (\epsilon \rightarrow -\epsilon) \end{aligned}$$

and

$$\begin{aligned} g(x) &= -2\pi \int_\theta \frac{1}{\omega/m} I_1(0, \theta, -\theta, 0) \frac{1}{\omega/m - x} = -\frac{1}{2} \int_{-\infty}^{\infty} d\theta \left( \frac{\coth^4 \frac{1}{2}(\theta - i\epsilon) \tanh^2 \theta + (\epsilon \rightarrow -\epsilon)}{\cosh \theta (\cosh \theta - x)} \right) \\ &= \frac{16}{1-x} - \frac{15\pi}{2x} - \frac{8}{x} - \frac{4\pi + 2}{x^2} - \frac{\pi}{x^3} - \frac{(x+1)^2 \sqrt{x^2 - 1}}{x^3(x-1)^2} - 2 \ln(-x + \sqrt{x^2 - 1}) \end{aligned} \quad (B1)$$

with  $g(0) = 10\pi + \frac{94}{3}$  (see also [21]) and the imaginary part for  $x > 1$ ,

$$\text{Im } g(x \pm i\epsilon) = \pm \Theta(x - 1) 2\pi \frac{(x+1)^2 \sqrt{x^2 - 1}}{x^3(x-1)^2}. \quad (B2)$$

**3. The four-point  $\Xi$  function and calculation of  $\chi(\omega_i, \omega_f)$**

The sum over all permutations in (20) yields

$$\begin{aligned} \Xi(k) = & \frac{1}{32\pi m^6} \left\{ \left( \frac{m}{k_4^0 + m} + \frac{m}{k_3^0 + m} \right) \left( \frac{m}{-k_1^0 + m} + \frac{m}{-k_2^0 + m} \right) g \left( \frac{-1}{2m} (k_3^0 + k_4^0) \right) \right. \\ & + \left( \frac{m}{k_4^0 + m} + \frac{m}{k_2^0 + m} \right) \left( \frac{m}{-k_1^0 + m} + \frac{m}{-k_3^0 + m} \right) g \left( \frac{-1}{2m} (k_2^0 + k_4^0) \right) \\ & + \left( \frac{m}{k_4^0 + m} + \frac{m}{k_1^0 + m} \right) \left( \frac{m}{-k_2^0 + m} + \frac{m}{-k_3^0 + m} \right) g \left( \frac{-1}{2m} (k_1^0 + k_4^0) \right) \\ & + \left( \frac{m}{k_3^0 + m} + \frac{m}{k_1^0 + m} \right) \left( \frac{m}{-k_2^0 + m} + \frac{m}{-k_4^0 + m} \right) g \left( \frac{-1}{2m} (k_1^0 + k_3^0) \right) \\ & + \left( \frac{m}{k_2^0 + m} + \frac{m}{k_1^0 + m} \right) \left( \frac{m}{-k_3^0 + m} + \frac{m}{-k_4^0 + m} \right) g \left( \frac{-1}{2m} (k_1^0 + k_2^0) \right) \\ & \left. + \left( \frac{m}{k_2^0 + m} + \frac{m}{k_3^0 + m} \right) \left( \frac{m}{-k_1^0 + m} + \frac{m}{-k_4^0 + m} \right) g \left( \frac{-1}{2m} (k_3^0 + k_2^0) \right) \right\}. \end{aligned}$$

Substituting this into (6) we obtain

$$\begin{aligned} \chi(\omega_i, \omega_f) \sim & \frac{(\omega_i - \omega_f + 2m)^2 \operatorname{Im} g \left[ \frac{-1}{2m} (\omega_i - \omega_f - i\delta_{12}) \right]}{(\omega_i + m)^2 (\omega_f - m)^2} + \frac{(\omega_i + \omega_f + 2m)^2 \operatorname{Im} g \left[ \frac{-1}{2m} (\omega_i + \omega_f - i\delta_{12}) \right]}{(\omega_i + m)^2 (\omega_f + m)^2} \\ & + \frac{(\omega_i + \omega_f - 2m)^2 \operatorname{Im} g \left[ \frac{1}{2m} (\omega_i + \omega_f + i\delta_{12}) \right]}{(\omega_f - m)^2 (\omega_i - m)^2} + \frac{(\omega_i - \omega_f - 2m)^2 \operatorname{Im} g \left[ \frac{1}{2m} (\omega_i - \omega_f + i\delta_{12}) \right]}{(\omega_f + m)^2 (\omega_i + m)^2}. \end{aligned}$$

At  $\omega_i > \omega_f > 0$  only the last two terms remain and (8) follows with  $G(x) = (x - 1)^2 \operatorname{Im} g(x)$ .

**APPENDIX C: PROOF OF (4)**

We consider the three-point Green's function

$$\tau_{\varphi\varphi\epsilon}(\underline{x}) = \langle 0 | T \varphi(x_1) \varphi(x_2) \epsilon(x_3) | 0 \rangle$$

and the Fourier transform (for  $\varphi_1, \varphi_2, \varphi_3 = \varphi, \varphi, \epsilon$ )

$$\begin{aligned} \tilde{\tau}_{\varphi\varphi\epsilon}(\underline{k}) &= \int \underline{d}^2x e^{i x_i k_i} \tau_{\varphi_1 \varphi_2 \varphi_3}(\underline{x}) = \sum_{\pi \in \mathcal{S}_3} \int \underline{d}^2x e^{i x_i \pi k_i} \Theta_{123} \langle \varphi_{\pi 1}(x_1) \varphi_{\pi 2}(x_2) \varphi_{\pi 3}(x_3) \rangle \\ &= (2\pi)^2 \delta^{(2)}(k_1 + k_2 + k_3) \tilde{\Xi}_{\varphi\varphi\epsilon}(\underline{k}). \end{aligned} \tag{C1}$$

The three-point Wightman functions in low intermediate particle number approximation are

$$\begin{aligned} w_{\varphi\varphi\epsilon}^{12}(\underline{x}) &= \frac{1}{2!} \int_{\underline{\theta}} \langle 0 | \varphi(x_1) | \theta_1 \rangle \langle \theta_1 | \varphi(x_2) | \theta_2, \theta_3 \rangle \langle \theta_3, \theta_2 | \epsilon(x_3) | 0 \rangle, \\ w_{\varphi\epsilon\varphi}^{11}(\underline{x}) &= \int_{\underline{\theta}} \langle 0 | \varphi(x_1) | \theta_1 \rangle \langle \theta_1 | \epsilon(x_2) | \theta_2 \rangle \langle \theta_2 | \varphi(x_3) | 0 \rangle, \\ w_{\epsilon\varphi\varphi}^{21}(\underline{x}) &= \frac{1}{2!} \int_{\underline{\theta}} \langle 0 | \epsilon(x_1) | \theta_1, \theta_2 \rangle \langle \theta_1, \theta_2 | \varphi(x_2) | \theta_3 \rangle \langle \theta_3 | \varphi(x_3) | 0 \rangle. \end{aligned}$$

As above using the form factor formulas (23) and (22) and the crossing relations (A1) one obtains the Fourier transforms [for  $k_i = (k_i^0, 0)$ ]

$$\begin{aligned} \tilde{\Xi}_{\varphi\varphi\epsilon}^{12}(k_1, k_2, k_3) &= \frac{-i}{32\pi m^4} \frac{m}{k_1^0 - m + i\epsilon} h \left[ -k_3^0 / (2m) + i\epsilon \right], \\ \tilde{\Xi}_{\varphi\epsilon\varphi}^{11}(k_1, k_2, k_3) &= -\frac{1}{4} \frac{i}{m^4} \frac{m}{m - k_1^0 - i\epsilon} \frac{m}{m + k_3^0 - i\epsilon}, \\ \tilde{\Xi}_{\epsilon\varphi\varphi}^{21}(k_1, k_2, k_3) &= \frac{-i}{32\pi m^4} \frac{m}{-k_3^0 - m + i\epsilon} h \left[ k_1^0 / (2m) + i\epsilon \right] \end{aligned}$$



with

$$h(x) = \int_{-\infty}^{\infty} \left(1 + \frac{1}{\cosh \theta}\right)^2 \frac{1}{\cosh \theta - x} d\theta = -\frac{2}{x}\pi - \frac{2}{x} - \frac{1}{x^2}\pi - 2 \frac{x^2 + 2x + 1}{x^2 \sqrt{x^2 - 1}} \ln(-x - \sqrt{x^2 - 1}).$$

Finally with (C1) we obtain

$$\tilde{\Xi}_{\varphi\varphi\epsilon}(k_1, k_2, k_3) = \tilde{\Xi}_{\varphi\varphi\epsilon}^{12}(k_1, k_2, k_3) + \tilde{\Xi}_{\varphi\varphi\epsilon}^{11}(k_1, k_3, k_2) + \tilde{\Xi}_{\epsilon\varphi\varphi}^{21}(k_3, k_1, k_2) + (k_1 \leftrightarrow k_2)$$

which proves (4).

- [1] Y. R. Shen, *The Principles of Nonlinear Optics* (Wiley, New York, 1984).
- [2] D. C. Hanna, M. A. Yuratic, and D. Cotter, *Nonlinear Optics of Free Atoms and Molecules* (Springer-Verlag, Berlin, 1979).
- [3] B. M. McCoy and T. T. Wu, *The Two-Dimensional Ising Model* (Dover, New York, 2014).
- [4] A. M. Tselik, *Quantum Fields Theory in Condensed Matter Physics*, 2nd ed. (Cambridge University Press, Cambridge, 2003), Chap. 28.
- [5] V. P. Yurov and A. B. Zamolodchikov, *Int. J. Mod. Phys. A* **6**, 3419 (1991).
- [6] R. Coldea, D. A. Tennant, E. M. Wheeler, E. Wawrzynska, D. Prabhakaran, M. Telling, K. Habicht, P. Smeibidl, and K. Kiefer, *Science* **327**, 177 (2010).
- [7] T. Liang, S. M. Koohpayeh, J. W. Krizan, T. M. McQueen, R. J. Cava, and N. P. Ong, *Nat. Commun.* **6**, 7611 (2015).
- [8] C. M. Morris, R. Valdés Aguilar, A. Ghosh, S. M. Koohpayeh, J. Krizan, R. J. Cava, O. Tchernyshyov, T. M. McQueen, and N. P. Armitage, *Phys. Rev. Lett.* **112**, 137403 (2014).
- [9] H. A. Kramers and W. Heisenberg, *Z. Phys.* **31**, 681 (1925).
- [10] P. A. M. Dirac, *Proc. R. Soc. London* **114**, 710 (1927).
- [11] B. S. Shastri and B. I. Shraiman, *Phys. Rev. Lett.* **65**, 1068 (1990); *Int. J. Mod. Phys. B* **5**, 365 (1991).
- [12] A. M. Shvaika, O. Vorobyov, J. K. Freericks, and T. P. Devereaux, *Phys. Rev. B* **71**, 045120 (2005).
- [13] S. T. Cundiff and S. Mukamel, *Phys. Today* **66**(7), 44 (2013).
- [14] O. Roslyak and S. Mukamel, *Phys. Rev. A* **79**, 063409 (2009).
- [15] M. Nakano, R. Kishi, S. Ohta, H. Takahashi, T. Kubo, K. Kamada, K. Ohta, E. Botek, and B. Champagne, *Phys. Rev. Lett.* **99**, 033001 (2007).
- [16] M. Karowski and H. J. Thun, *Nucl. Phys. B* **130**, 295 (1977).
- [17] H. Babujian, A. Foerster, and M. Karowski, *Nucl. Phys. B* **736**, 169 (2006).
- [18] B. Berg, M. Karowski, and P. Weisz, *Phys. Rev. D* **19**, 2477 (1979).
- [19] M. Karowski, in *Field Theoretic Methods in Particle Physics*, edited by W. Rühl (Plenum, New York, 1980), pp. 307–324, presented at Kaiserslautern NATO Inst., 1979.
- [20] W. G. Yin, X. Liu, A. M. Tselik, M. P. M. Dean, M. H. Upton, J. Kim, D. Casa, A. Said, T. Gog, T. F. Qi, G. Cao, and J. P. Hill, *Phys. Rev. Lett.* **111**, 057202 (2013).
- [21] J. Balog, M. Niedermaier, F. Niedermayer, A. Patrascioiu, E. Seiler, and P. Weisz, *Nucl. Phys. B* **583**, 614 (2000).
- [22] H. Babujian and M. Karowski, *Nucl. Phys. B* **620**, 407 (2002).